

# Medical Progress

## Prostaglandins, Thromboxanes and Leukotrienes in Clinical Medicine

ROBERT D. ZIPSER, MD, *Torrance, California*, and GIACOMO LAFFI, MD, *Florence, Italy*

*Although prostaglandin research began about 50 years ago, many of the most important advances in understanding the biochemistry, physiology and pharmacology have taken place within the past five to ten years. There is great potential for the extension of this research to the clinical practice of medicine. At this time, the most common interaction that clinicians have with the prostaglandin field is in administering nonsteroidal anti-inflammatory drugs, which function by inhibiting prostaglandins. The uses of these drugs include treating not only inflammation, but also dysmenorrhea, some renal diseases, thrombotic diseases and some metabolic disorders. Prostaglandin analogs, with their potent effects on uterine contraction, are in common use in obstetrics. Other analogs with gastric and duodenal cytoprotective effects are useful in treating peptic ulcer disease. Future benefits from prostaglandin and leukotriene research may include new therapy for inflammatory and hypersensitivity diseases such as asthma, inflammatory bowel disease and dermatitis.*

(Zipser RD, Laffi G: Prostaglandins, thromboxanes and leukotrienes in clinical medicine [Medical Progress]. West J Med 1985 Oct; 143:485-497)

In this review we emphasize those areas of prostaglandin, thromboxane and leukotriene (collectively called "eicosanoids") biochemistry and physiology that interact with the contemporary practice of medicine. Additional emphasis will include discussions of assay methods, inhibitors and pharmacology to assist readers in interpreting the increasing number of publications that cover these areas. Eicosanoid research has made important contributions to several subspecialties, including cardiovascular and thrombotic disease, immunity and inflammation, reproduction, nephrology, pulmonary disease, gastroenterology and metabolic disorders. These areas will be discussed in more detail.

### History and Introduction

The history of prostaglandins began in the 1930s with the observation by two New York gynecologists, Kurzrok and Lieb, that human semen caused contractions and relaxation of human myometrium.<sup>1</sup> These observations were soon confirmed by Goldblatt in England<sup>2</sup> and by von Euler in Sweden.<sup>3</sup> In 1935 von Euler purified the active compounds by acid lipid extraction of sheep vesicular glands and coined the term "prostaglandin" in the belief that the prostate was the source.<sup>4</sup> At about the same time, the history of leukotrienes began with the finding by Feldberg, Kellaway and Trethewie

of a factor from dog lung exposed to cobra venom and a similar factor from sensitized guinea pig lung that caused a slow, prolonged contraction of the guinea pig jejunum.<sup>5,6</sup> These factors were called slow-reacting substance of anaphylaxis (SRS-A) and later identified as a possible mediator of asthma and other hypersensitivity reactions. There was then a delay in progress until the 1960s when a series of major advances began in Sweden by Bergström and colleagues, including Samuelsson, and in The Netherlands by Van Dorp. Prostaglandins were identified as a group of compounds rather than a single substance, and arachidonic acid was identified as their precursor.<sup>7,8</sup> Isolation of large amounts of natural prostaglandins from the Caribbean coral *Plexaura homomalla*, new methods of synthesis pioneered by Corey and others and extensive distribution of these products to investigators by the Upjohn Company and other pharmaceutical companies permitted intensive widespread study of these compounds. It was soon recognized that prostaglandins were produced by nearly all biologic tissue. Profound pharmacologic effects of these compounds were shown on smooth muscle contraction, cell secretions and platelet aggregation, which fostered the concept that prostaglandins functioned as local mediators of many biologic systems.

In 1971 the number of publications in this field began to


From the Division of Gastroenterology, Department of Medicine, Harbor-UCLA Medical Center, Torrance, California. Dr Laffi was a visiting scholar from Istituto di Clinica Medica IV, Università degli Studi di Firenze, Italy, when this study was done.

This study was supported by National Institutes of Health grant No. AM-33410.

Reprint requests to Robert D. Zipser, MD, Gastroenterology, C-1 Trailer, Harbor-UCLA Medical Center, 1000 W Carson St, Torrance, CA 90509.

# ABBREVIATIONS USED IN TEXT

cyclic AMP = adenosine 3':5'-cyclic phosphate  
HETE = hydroxyeicosatetraenoic acid  
HPETE = hydroperoxyeicosatetraenoic acid  
LT = leukotriene  
NSAID = nonsteroidal anti-inflammatory drug  
PG = prostaglandin  
SRS-A = slow-reacting substance of anaphylaxis  
Tx = thromboxane



**Figure 1.**—Fatty acid precursors of eicosanoids.

explode with the key discovery by Vane and colleagues in England that aspirin, indomethacin and other nonsteroidal anti-inflammatory drugs inhibited prostaglandin biosynthesis.<sup>9</sup> Over the past 12 years there have been more than 26,000 publications. Major highlights included the 1975 report by Hamberg, Svensson and Samuelsson identifying the potent platelet-aggregatory substance, thromboxane  $A_2$ ,<sup>10</sup> and the 1976 identification by Moncada, Vane and co-workers of the antiaggregatory vasodilator, prostaglandin  $I_2$ .<sup>11</sup> During this time, parallel progress was made in studies of SRS-A including identification of arachidonic acid as the precursor.<sup>12</sup> In 1979 the Samuelsson laboratory announced the structures of the leukotrienes that make up the compounds of SRS-A.<sup>13,14</sup> New enzymatic pathways of arachidonic acid metabolism involving the lipoxygenase pathway were identified, as well as potent new biologic mediators of inflammation. In 1982 the Nobel prize for physiology and medicine was awarded to Sune Bergström, Bengt Samuelsson and John Vane for their major contributions to this field. The field continues to expand with an escalating number of publications on new metabolites of arachidonic acid, new inhibitors and a plethora of data on prostaglandin involvement in physiology.

## Biochemistry

Prostaglandins and thromboxanes are derived from 20-carbon fatty acids. In mammals, arachidonic acid is the major precursor of prostaglandins ("2" series prostaglandins) and is localized primarily in the membrane phospholipids. Dihomo- $\gamma$ -linolenic acid is the precursor of the "1" series prostaglandins (present only in trace amounts), and eicosapentaenoic acid (predominant in fish oils) is converted to some extent to the "3" series prostaglandins (Figure 1). Extreme variations in dietary fat and eicosapentaenoic acid, such as that derived from the marine diet of Greenland Eskimos, can compete with the enzymes governing arachidonic acid metabolism and alter prostaglandin and thromboxane synthesis, resulting in biologic changes such as prolonged bleeding time.<sup>15</sup> In essential fatty acid deficiency, prostaglandin synthesis is diminished, which may account for some of the clinical manifestations such as dermatopathy.

The cascade of prostaglandin synthesis begins with the release of arachidonic acid from phospholipids via activation of phospholipase (primarily phospholipase  $A_2$ ). This rate-limiting step is initiated by hormonal, ischemic, neural, inflammatory or other stimuli varying from cell to cell. Once released, arachidonic acid is available for enzymatic oxidation into hydroperoxy and hydroxy derivatives. For example, the 12-lipoxygenase enzyme in platelets induces formation of the unstable 12-hydroperoxyeicosatetraenoic acid (12-HPETE), which is reduced by peroxidase to the stable 12-hydroxyeicosatetraenoic acid (12-HETE), with unclear bio-

logic activity. By a similar mechanism, 15-lipoxygenase in leukocytes forms 15-HPETE and 15-HETE, which may have some inflammatory activity. Two other enzymatic oxidation sites at carbon 11 and 5 are of more biologic importance, leading to formation of prostaglandins and leukotrienes, respectively. Aside from the biologic activity of the arachidonic acid metabolites, the oxidation-peroxidation reactions may have more direct biologic effects by forming oxygen radicals, which may induce tissue injury and carcinogenesis.<sup>16,17</sup>

Oxidation in the microsomes of most mammalian cells by the cyclooxygenase enzyme (originally called prostaglandin synthetase) at C 11 induces formation of a cyclopentane ring and a cyclic endoperoxide (prostaglandin [PG]  $G_2$ ) (Figure 2).  $PGG_2$  is rapidly converted by peroxidase activity to  $PGH_2$ , the unstable common intermediate of prostaglandins and thromboxanes. Conversion of  $PGH_2$  to thromboxanes or to various prostaglandins is effected by enzyme systems characteristic for each specific cell type. Several examples are listed.  $PGE_2$  and  $PGF_{2\alpha}$  are the classic prostaglandins present in many tissues, including the kidney and uterus, with diverse possible physiologic actions including smooth muscle contraction, pyretic reactions and inflammatory interactions.  $PGI_2$  (prostacyclin) is particularly prominent in vascular endothelium and possesses both vasodilatory and antiaggregatory actions.  $PGD_2$  is a major product of mast cells and brain tissue. Thromboxane  $A_2$  production is most commonly associated with platelet aggregation and vasoconstriction (see reviews.)<sup>18-20</sup>

Leukotriene formation, which predominates in polynuclear leukocytes and some mononuclear cells, results from 5-lipoxygenase activity forming 5-HPETE and conversion to an epoxide moiety, the unstable common intermediate, leukotriene (LT)  $A_4$ .  $LTA_4$  can then react with water to form the potent chemokinetic and chemotactic leukotriene,  $LTB_4$ . Alternatively,  $LTA_4$  reacting with glutathione leads to the peptide derivatives,  $LTC_4$ ,  $LTD_4$  and  $LTE_4$ , which make up the potent inflammatory mediators collectively known as SRS-A (see reviews.)<sup>12,21-24</sup> Recently, new compounds and pathways have been proposed, such as metabolites of 15-lipoxygenase, called lipoxins that may also have inflammatory functions.<sup>25</sup>

## Eicosanoid Inhibitors


By far the most common interaction that clinicians have with the field of prostaglandins is the therapeutic use of nonsteroidal anti-inflammatory drugs (NSAIDs) and of corticosteroids for inflammatory conditions. The use of anti-inflammatory agents, however, certainly preceded prostaglandin research (willow bark has been used for centuries,

for instance, and acetylsalicylic acid has been commercially available since about 1900), and it is not established that all the effects of these drugs can be explained by their inhibiting the cyclooxygenase enzyme. A detailed review of NSAID pharmacology and clinical experience exceeds the scope of this summary, and excellent reviews are available.<sup>26-32</sup> Instead, in this summary we will briefly consider the modes of action of these agents, the prospect for more specific inhibitors and the relevance of the effects of these agents to our understanding of prostaglandin physiology.

In general, NSAIDs function by inhibiting the cyclooxygenase enzyme, thereby diminishing the conversion of arachidonic acid to  $\text{PGH}_2$  and to other prostaglandins and thromboxanes.<sup>9,13</sup> The anti-inflammatory, analgesic and antipyretic actions of aspirinlike drugs are attributed to inhibition of prostaglandins, particularly  $\text{PGE}_2$ . Other anti-inflammatory drugs such as colchicine and gold do not inhibit prostaglandin. The NSAID effects on cyclooxygenase include diminished peroxidase activity and reduced generation of oxygen radicals, which may also reduce tissue damage and possibly reduce the formation of carcinogens. The anti-inflammatory potency of NSAIDs generally parallels the prostaglandin-inhibiting potency, although there is some variation in different tissues. For example, compared with newer

NSAIDs, aspirin is a relatively weak prostaglandin inhibitor, but it causes a more prolonged inhibition of platelet thromboxane because the irreversibly acetylated cyclooxygenase enzyme cannot be regenerated in the life span of these cells. Acetaminophen is a poor cyclooxygenase inhibitor in most in vitro systems, but there is evidence that it may be effective in brain tissue, explaining its antipyretic and analgesic effects without peripheral anti-inflammatory effects.<sup>32</sup> Compared with indomethacin and ibuprofen, the new NSAID sulindac is less likely to cause acute renal impairment in patients with chronic glomerulonephritis or in those with cirrhosis and ascites<sup>33,34</sup>; however, it is not clear if this apparent renal sparing of sulindac is the result of rapid inactivation of the drug by the kidney or of overall weaker cyclooxygenase activity. Sodium salicylate reduces prostaglandin synthesis in sites of inflammation and in the pancreatic islet without inhibiting gastric mucosal or platelet prostaglandins.<sup>35</sup>

Finally, caution is urged in extrapolating specific physiologic functions of prostaglandins from the effects of NSAIDs. The NSAIDs are considered "nonspecific" because they inhibit all of the cyclooxygenase products and also because they act on multiple organ systems. The observed effects may be the culmination of effects on several biologic systems. These agents also have nonprostaglandin effects. For example, indo-



**Figure 2.**—Eicosanoid biochemical pathways. HETE = hydroxyeicosatetraenoic acid; HPETE = hydroperoxyeicosatetraenoic acid; LT ( $A_4$ ,  $B_4$ ,  $C_4$ ,  $D_4$ ,  $E_4$ ) = leukotriene; NSAIDs = nonsteroidal anti-inflammatory drugs; PG ( $D_2$ ,  $E_2$ ,  $F_{1\alpha}$ ,  $F_{2\alpha}$ ,  $G_2$ ,  $H_2$ ,  $I_2$ ) = prostaglandin; Tx ( $A_2$ ,  $B_2$ ) = thromboxane

methacin in high concentrations affects protein kinase, calcium flux and other enzymes.<sup>29</sup> In addition, at least some of the NSAIDs also block arachidonic acid metabolism by the 15-lipoxygenase enzyme, perhaps contributing to the anti-inflammatory effects.<sup>30</sup>

Inhibitors of specific enzyme pathways distal to the cyclooxygenase enzyme have recently been developed with possible clinical implications. The first to undergo clinical trials have been imidazole and pyridine derivatives, which selectively block thromboxane synthetase without reducing other cyclooxygenase products.<sup>36,37</sup> The premise is that these drugs will have antithrombotic action because they block platelet thromboxane without reducing blood vessel synthesis of prostacyclin. Early clinical trials of patients with thrombotic and vasospastic disease have suggested some benefit but no clear advantage over aspirin. Clinical application may have more promise in other disease states such as glomerulonephritis.<sup>38</sup>

Pharmacologic administration of glucocorticoids causes many effects unrelated to arachidonic acid metabolism. At least in some biologic systems, these steroids inhibit the release of arachidonic acid from membrane phospholipids.<sup>39</sup> This effect involves synthesis of a glycoprotein that inhibits phospholipase A<sub>2</sub> activity.<sup>40,41</sup> The new term "lipocortins" best describes this family of compounds.<sup>42</sup> In susceptible tissue, all arachidonic acid metabolites are inhibited, including cyclooxygenase and lipoxygenase systems. The anticipated reduction in leukotriene synthesis is the popular explanation for some of the glucocorticoid efficacy in cases of hypersensitivity reaction such as asthma and in inflammatory states. Fostered by these studies and the increasing amount of data on the potent effects of leukotrienes, there is currently an intense effort to develop specific leukotriene inhibitors for clinical uses that may include patients with asthma, inflammatory arthritis, dermatitis and inflammatory bowel disease.<sup>43</sup>

### Methodologic Constraints

It has been the unfortunate recurring history of prostaglandin assays in clinical research that the early studies of each eicosanoid report biologic levels that greatly exceed those subsequently determined with better methods. The physiologic roles equated with the early measurements are then subsequently disproved. For example, many early studies of plasma and serum PGE<sub>1</sub>, PGE<sub>2</sub>, PGA<sub>1</sub> and PGA<sub>2</sub> reported normal values ranging from several hundred picograms to several hundred nanograms per milliliter. Theories on the pathogenesis of blood pressure regulation, sodium balance and diarrheal states were widely promulgated based on these reports. Later studies using mass spectrometry confirmed that PGA<sub>1</sub> and PGA<sub>2</sub> did not exist in detectable amounts in human circulation and that PGE<sub>2</sub> plasma levels were less than 30 pg per ml. Similar misadventures have occurred with reports of circulating levels of prostacyclin (PGI<sub>2</sub>), thromboxane B<sub>2</sub>, PGF<sub>2α</sub> and some prostaglandin metabolites (see reviews).<sup>19,44,45</sup> It is now clear that none of these compounds are present in sufficient concentration to act as circulating hormones; instead, eicosanoids function primarily as local mediators. The methodologic problems are caused by several factors, which include poor antiserum specificity in early radioimmunoassays, the myriad of unidentified prostaglandin metabolites that may cross-react in these assays, poor specificity and sensitivity of bioassays, the pronounced poten-

tial for artifactual generation of eicosanoids during sample collection—such as from platelet release and vessel wall trauma during blood sampling—the short biologic and chemical half-life of many eicosanoids and unidentified interfering substances in early mass spectrometry. It behooves the reader of prostaglandin research to be cautious of all reports of circulating blood and urine levels unless extensive validating studies (particularly comparison with mass spectrometry) have been presented. Measurements of tissue concentration of eicosanoids are suspect because these compounds are generally not stored and because they are readily produced by tissue trauma, such as by biopsy. One technique to avoid these problems is to measure stable metabolites. For example, urinary excretion of 2,3-dinor-PGF<sub>1α</sub> and 2,3-dinor-thromboxane B<sub>2</sub> appear to quantitatively reflect systemic production of prostacyclin and thromboxane, respectively,<sup>46</sup> and plasma assay of a tetranor metabolite of PGF<sub>2α</sub>, which has a long half-life in circulation, accurately reflects systemic production of PGF<sub>2α</sub>.<sup>47</sup>

### Cardiovascular and Thrombotic Diseases

The role of prostaglandins in controlling blood pressure has been a topic of dispute for almost 20 years since the isolation of prostaglandin A<sub>2</sub> from kidney medulla and the early reports claiming that circulating levels of PGA<sub>2</sub> were of sufficient concentration to affect blood pressure.<sup>48–50</sup> Later studies showed that PGA<sub>2</sub> was not present in the circulation.<sup>51</sup> Similarly, other vasodepressor prostaglandins such as PGI<sub>2</sub> exist in blood in concentrations too low to have circulating hormonal effects.<sup>52</sup> Nevertheless, it is likely that local vascular production of vasodilatory prostaglandins and possibly vasoconstrictor thromboxanes do function to modulate local vascular resistance, especially in response to vasopressor hormones and to volume. Thus, administering NSAIDs may increase blood pressure (aside from the sodium-retaining effects) in a few patients such as those with mineralocorticoid-induced hypertension,<sup>53</sup> and NSAIDs may blunt the effects of antihypertensive drugs.<sup>54</sup> The hemodynamic effects of these agents are particularly evident in patients with congestive heart failure.<sup>55</sup>

A more contemporary issue is the possible role of prostacyclin (PGI<sub>2</sub>) and thromboxane (Tx) A<sub>2</sub> in regulating platelet aggregation in patients with thrombotic and atherosclerotic disease. Thromboxane A<sub>2</sub>, a major arachidonic acid product of platelets, is a potent vasoconstrictor (similar in potency to angiotensin II), platelet-aggregatory stimulus (independent of adenosine diphosphate and thrombin) and bronchoconstrictor.<sup>10</sup> TxA<sub>2</sub> has a short half-life (30 seconds in aqueous media) and spontaneously converts to the inactive TxB<sub>2</sub>. Following vascular wall trauma, collagen and other factors stimulate platelet aggregation and thromboxane release. Aspirin, other NSAIDs and selective thromboxane-synthesis inhibitors block thromboxane generation and prolong bleeding time in vivo and platelet aggregation in vitro.<sup>9,56–58</sup> Opposing effects result from PGI<sub>2</sub>, the predominant arachidonic acid metabolite of intact vascular endothelium with vasodilatory and bronchodilatory properties.<sup>11</sup> PGI<sub>2</sub> inhibits platelet aggregation and prevents TxA<sub>2</sub> production via a platelet adenosine 3':5'-cyclic phosphate (cyclic AMP)-mediated pathway.<sup>57,59</sup> Dipyridamole, an inhibitor of phosphodiesterase, also increases cyclic-AMP levels, inhibits platelet aggregation by cyclic

AMP of other mechanisms and potentiates the antiaggregatory effect of aspirin.<sup>60</sup> As with  $\text{TxA}_2$ ,  $\text{PGI}_2$  has a short half-life (about 1.5 minutes in aqueous solution), and it is converted to an inactive metabolite, 6-keto-prostaglandin  $\text{F}_{1\alpha}$ , but possibly also to an active metabolite, 6-keto-PGE<sub>1</sub>, in some tissues. The general interpretation of these studies is that the intact endothelium prevents platelet adherence and aggregation via local generation of  $\text{PGI}_2$  (plus several other antiplatelet factors). With disruption of the endothelium, initial platelet adherence occurs and platelet  $\text{TxA}_2$  is then produced that causes local vasoconstriction and augments platelet aggregation at the injury site. Production of  $\text{PGI}_2$  from normal endothelium adjacent to the injury site may be increased by the use of endoperoxide precursors ( $\text{PGH}_2$ ) released during arachidonic acid metabolism from the platelets and taken up by the endothelium, thus preventing spread of platelet aggregation to normal endothelium (Figure 3).<sup>57,59-61</sup>


One clinical implication of these opposing effects of  $\text{TxA}_2$  and  $\text{PGI}_2$  is the concept that an imbalance of these products may contribute to thrombotic, atherosclerotic and vasospastic disease.<sup>62</sup> This concept is supported by several in vitro studies showing increased thromboxane generation from disease states susceptible to atherosclerosis, such as hypercholesterolemia<sup>63</sup> and diabetes mellitus.<sup>64</sup> Conversely, high-density lipoproteins are associated with a reduced risk of cardiovascular disease and in vitro studies have shown an increased endothelial  $\text{PGI}_2$  response to high-density lipoprotein.<sup>65</sup> In addition, smoking decreases the urinary excretion of a  $\text{PGI}_2$  metabolite.<sup>66</sup> Although the theories of imbalances of arachidonic acid metabolites as a cause of atherosclerosis are intriguing, rigorous evidence of clinical relevance is still lacking. Indeed, recent data suggest that systemic  $\text{PGI}_2$  production may actually be increased rather than decreased in patients with atherosclerosis.<sup>67</sup> Based on this imbalance theory, therapeutic strategies to prevent cardiovascular disease that attempt to selectively reduce  $\text{TxA}_2$  without impairing  $\text{PGI}_2$  have been proposed, including administering very low doses of aspirin (to block platelet but not vascular cyclooxygenase), administering selective inhibitors of thromboxane synthetase and prescribing diets rich in eicosapenta-

noic acid (fish oils).<sup>68</sup> Although controversy persists, increasing data suggest that administering aspirin (with or without dipyridamole) is of benefit in reducing the risk of myocardial infarction in some patients and in reducing reocclusion of saphenous vein coronary artery bypass grafts.<sup>69-71</sup> These data provide clinical support at least for the importance of platelet aggregation and thromboxane in these diseases. A role for  $\text{PGI}_2$  is likely, but not proved.<sup>72</sup>

Another exciting extension of prostaglandin research to cardiovascular disease is the pharmacologic administration of prostacyclin ( $\text{PGI}_2$ ) or a stable analog for its platelet antiaggregatory and vasodilatory effects.<sup>73</sup> Prostacyclin has been substituted for heparin during extracorporeal circulation such as hemodialysis, cardiopulmonary bypass and charcoal perfusion.<sup>74-77</sup> Compared with the effects of heparin therapy, thrombocytopenia is reduced or prevented, most platelet functions are maintained and blood loss may also be reduced. The use of prostacyclin has also been advocated for treating patients who have atherosclerotic obstruction of extremities, central retinal vein occlusion, angina pectoris or ischemic strokes based on the premise that atherosclerotic arteries have decreased prostacyclin production. Initial results of uncontrolled studies are promising.<sup>78,79</sup> Earlier studies using infusions of PGE<sub>1</sub> also claimed relief of pain and ulcer healing in patients with chronic ischemia of the legs.<sup>80</sup> Open trials of prostacyclin infusion in patients with the Raynaud's phenomenon have also suggested prolonged improvement.<sup>81</sup> Results of double-blind controlled trials have suggested that these infusions are not consistently beneficial<sup>82-84</sup> and such therapy will probably not become clinically useful.

## Immune and Inflammatory Responses

Perhaps the areas with the most rapidly expanding knowledge about eicosanoid physiology are those of immunity and inflammation.<sup>21-23,28,31,85-88</sup> Increased production of arachidonic acid metabolites occurs at sites of inflammation,<sup>85-89</sup> arising from the injured tissue, from inflammatory cells and from noninflammatory cells that are apparently stimulated by other products of inflammation.<sup>90,91</sup> Oxygen free radicals are generated by both cyclooxygenase and lipoxygenase pathways, and these products likely contribute to tissue injury.<sup>92</sup> Several prostaglandins, especially PGE<sub>2</sub>,  $\text{PGI}_2$  and possibly  $\text{PGD}_2$ , contribute to inflammation by increasing the blood flow (erythema), by dilating vessels and increasing capillary permeability (edema) and by increasing pain sensitivity to other mediators such as to bradykinin and histamine. Prostaglandins also have pyretic effects under some conditions such as following release of leukocytic pyrogen and endotoxin.<sup>93,94</sup> It is unclear whether the cyclooxygenase products have chemotactic and chemokinetic properties: discrepant results have been reported in several studies. However, leukotriene B<sub>4</sub> is a very potent chemotactic factor for neutrophils, eosinophils and mononuclear cells.<sup>31,95</sup> Some mono-HETEs are less potent chemotactic agents. LTB<sub>4</sub> also activates neutrophils by causing degranulation and superoxide generation. LTC<sub>4</sub>, LTD<sub>4</sub> and LTE<sub>4</sub> (SRS-A) have vasoconstrictor activity; they also contribute to edema by increasing capillary permeability. These SRS-A factors are from 100-fold to 10,000-fold more potent than histamine in constricting pulmonary airways. It is apparent that the anti-inflammatory effects of NSAIDs and of corticosteroids function in part by inhibiting the eicosanoid



**Figure 3.**—Thromboxane (Tx) and prostacyclin (prostaglandin [ $\text{PGI}_2$ ]) interactions from platelets and blood vessel endothelium. cAMP = cyclic AMP (adenosine 3':5'-cyclic phosphate), NSAIDs = nonsteroidal anti-inflammatory drugs

actions cited above. It should be emphasized that eicosanoids act primarily as potentiators (or in some cases as modulators), rather than as initiators, of inflammation. In some inflammatory conditions the effects of certain eicosanoids may predominate, favoring the use of specific blockers of the arachidonic acid cascade. For example, lipoxygenase products may be particularly important in cases of psoriasis,<sup>96,97</sup> inflammatory bowel disease<sup>98</sup> and rheumatoid arthritis,<sup>99</sup> whereas PGD<sub>2</sub> predominates in mastocytosis.<sup>100</sup>

In contrast to the apparent proinflammatory actions, eicosanoids generally suppress immune functions.<sup>87,88</sup> In low concentrations, PGE<sub>2</sub> suppresses T-lymphocyte functions that include antigenic stimulation, lymphokine production, lymphocyte migration, mitogen responsiveness and clonal proliferation. These actions are mediated by enhanced generation of cyclic AMP. LTB<sub>4</sub> and 15-HPETE may have similar effects. NK-cell and monocyte cytolytic activity is inhibited by PGE<sub>2</sub> and by lipoxins, although lipoxins may act independently of cyclic AMP. Prostaglandins may also inhibit B-cell function, either directly or via actions on other immune cells. Macrophages and monocytes produce large amounts of PGE<sub>2</sub> in response to lymphokines and other stimuli that include zymosan, endotoxins, antigen-antibody complexes and immunoglobulin-G fragments. A popular theory is that antigens stimulate lymphocyte production of lymphokines, which activate macrophages to produce prostaglandins. These prostaglandins then function to inhibit the lymphocyte function.<sup>101</sup> In cases of Hodgkin's disease the defect in cellular immunity is due to a population of prostaglandin-producing suppressor cells.<sup>102</sup> Immunosuppressive activity from macrophages (presumably via prostaglandin production) may contribute to the anergy present in patients with chronic inflammatory diseases such as Q fever, coccidioidomycosis and tuberculosis.<sup>103</sup>

These pieces of information are consistent with the theory that eicosanoids function predominantly as potentiators of acute inflammation, actions that are inhibited by corticosteroids and by NSAIDs. In contrast, in some chronic disease states, eicosanoids function to depress cellular immunity and prostaglandin inhibitors may act to stimulate cellular immune function.

## Reproduction

Another important impact of prostaglandin research on clinical medicine involves reproduction, including the extensive use of exogenous prostaglandins as pharmacologic agents.<sup>104-107</sup>

Seminal fluid contains the highest concentration of prostaglandins of all human tissue, more than 100,000-fold more concentrated than in circulating plasma.<sup>108,109</sup> This observation has led to speculation that seminal fluid prostaglandins are involved in the contractile response of the vas deferens and the uterus, although the precise function in fertility is not established. There is some evidence, however, of diminished seminal fluid prostaglandin concentration in some infertile men who have no other apparent abnormality.<sup>110</sup>

There is pronounced species variation in the physiologic functions and pharmacologic effects of prostaglandins in female reproductive organs. In rats and rabbits, but not in humans, prostaglandins are obligatory mediators of ovulation.<sup>111,112</sup> Gonadotropin-stimulated cyclic AMP in the granulosa cell induces prostaglandin synthesis.<sup>107,113</sup> Indo-

methacin prevents follicular rupture, and exogenous prostaglandins reverse this effect. In many species including sheep, cows, horses and pigs, prostaglandin production (possibly PGF<sub>2α</sub> from the uterus) is the key factor causing luteolysis.<sup>113-115</sup> Prostaglandin administration has been used to synchronize the estrous cycle of cows and other animals to facilitate artificial insemination.<sup>116</sup> In humans, there is no strong evidence that prostaglandins or prostaglandin inhibitors have direct effects on ovulation or on maintaining the corpus luteum. There is substantial evidence, however, that increased production of endometrial prostaglandins contributes to the symptoms of dysmenorrhea, and NSAIDs are now widely prescribed for these symptoms.<sup>117-119</sup> Several clinical trials have confirmed the safety and efficacy of these drugs for such symptoms as uterine cramps, headache, nausea and diarrhea.<sup>120</sup> NSAIDs are also used to treat dysfunctional uterine bleeding, especially in association with intrauterine contraceptive devices.<sup>121,122</sup>

During pregnancy, the uterus is particularly sensitive to PGE<sub>2</sub>, PGF<sub>2α</sub> and their analogs. Endogenous prostaglandins may mediate uterine contractions in nonsurgical abortions,<sup>123</sup> and exogenous prostaglandins are widely used for terminating pregnancy, particularly in the second trimester.<sup>104,105</sup> Major complications of this treatment include bronchospasm, shock, seizures and cervicovaginal fistulas. Gastrointestinal symptoms are common. Prostaglandins are generally more effective, however, and are associated with less frequent serious side effects than hypertonic saline.<sup>124</sup> Side effects are also minimized with local administration (intrauterine or intravaginal instead of oral, intravenous or intramuscular) and with newer analogs.<sup>125,126</sup> Abnormal pregnancies, such as missed abortions and intrauterine fetal death, appear particularly sensitive to exogenous prostaglandins.<sup>127</sup>

Karim and co-workers in Singapore first reported the use of prostaglandins for inducing labor in 1968.<sup>128</sup> However, oxytocin generally remains the drug of choice.<sup>104,107,129</sup> NSAIDs may prolong gestation and delay the onset of labor. These drugs should not be used for treating cases of premature labor as they potentially affect fetal pulmonary circulation and hemostasis.<sup>130</sup> Endogenous prostaglandins may be involved in cervical ripening,<sup>131</sup> and exogenous prostaglandins have been used to hasten cervical ripening at term.<sup>132</sup> Prostaglandins may also be effective in controlling postpartum hemorrhage.

NSAIDs are an established therapy for closing a patent ductus arteriosus in premature newborns.<sup>133,134</sup> It is likely that prostaglandins (probably PGE<sub>2</sub> or PGI<sub>2</sub>) function to maintain patency of the ductus during normal fetal life. These prostaglandins arise from ductal tissue or possibly from the lung. At birth, the ductus normally closes in response to increased oxygen tension, possibly related to a fall or a shift in prostaglandin synthesis.<sup>135-137</sup> Exogenous prostaglandins may be infused to prevent closure of the ductus arteriosus when it is critical to maintain fetal circulation, such as with congenital right ventricular outflow obstruction, transposition of the great arteries and with some anomalies of the aortic arch.<sup>136</sup>

## Renal Function

There has been extensive research summarized by thorough reviews on the role of prostaglandins in renal physiology.<sup>138-144</sup> Prostaglandins have been implicated in the physio-



logic control of renal hemodynamics, renin release, sodium excretion and water transport. There are extensive data on possible functions of prostaglandins and thromboxanes in a number of renal diseases, including Bartter's syndrome, obstructive uropathy, functional renal impairment associated with liver disease, glomerulonephritis, acute and chronic renal failure and renal transplant rejection. In this review we will emphasize the clinical applications of these data, particularly the beneficial and adverse effects of NSAIDs.

Bartter's syndrome was the first renal disorder clearly recognized to involve augmented renal prostaglandin synthesis.<sup>145-147</sup> Increased renal synthesis of PGE<sub>2</sub> or PGI<sub>2</sub> from different regions of the kidney and peripheral vasculature contribute to the hyperreninism, sodium depletion, polyuria, low blood pressure, resistance to the pressor effects of angiotensin and norepinephrine, increased urinary excretion of prostaglandins and possibly to the hypokalemia. Administering indomethacin and other NSAIDs greatly alleviates these abnormalities, and NSAIDs are now the standard treatment. However, chloride wasting is likely the primary abnormality of this disorder and prostaglandins are probably secondary mediators. Giving NSAIDs does not fully correct the hypokalemia.<sup>148</sup> In patients with the more common disorders involving volume depletion or relative underperfusion of the kidneys, such as diuretic administration, surreptitious vomiting, congestive heart failure and cirrhosis with ascites, the increased prostaglandin excretion and prostaglandin-mediated hyperreninism are also reduced by NSAIDs.<sup>143, 144, 149-153</sup> Results of these clinical studies and of many animal and in vitro studies support the concept that prostaglandins (probably PGI<sub>2</sub> produced in the renal cortex) are involved in renin release, although prostaglandins are not obligatory intermediates, as profound stimuli arising from sodium depletion,  $\beta$ -adrenergic stimuli or hypotension causes renin release despite cyclooxygenase inhibition.<sup>154-158</sup> An additional clinical implication of these observations is that during renin determination for classification of hypertension or other conditions, NSAIDs will suppress renin activity and withdrawal of NSAIDs may cause a rebound elevation in renin.<sup>159</sup> Prolonged suppression of renin and the secondary suppression of aldosterone in some patients with mild chronic renal impairment may cause hyperkalemia and mimic the syndrome of hyporeninemic hypoaldosteronism.<sup>160</sup>

Synthesis of renal cortical vasodilatory prostaglandins is enhanced in patients with glomerulonephritis or underperfusion of the kidney, such as in those who have cirrhosis with ascites, heart failure or sodium depletion. These prostaglandins function to maintain renal perfusion, countering vasoconstrictor stimuli such as catecholamines and angiotensin. Administering NSAIDs to these susceptible patients unmasks the underlying renal vasoconstriction and results in an abrupt decrease in renal blood flow and glomerular filtration rate.\* This situation is most dramatic in patients with cirrhosis and ascites in whom as little as 25 mg of indomethacin may cause anuria for several hours.<sup>151-153</sup> A diminished glomerular filtration rate may contribute to the apparent beneficial effects of NSAIDs to reduce proteinuria in patients with nephrotic syndrome, although a more direct prostaglandin effect on proteinuria may also be involved.<sup>165</sup>

The role of renal prostaglandins in sodium excretion is incompletely understood. Clearly, various NSAIDs cause sodium retention—generally about 150 mEq over several days in healthy subjects with greater retention in patients with various disease states.<sup>166</sup> The intrarenal site of action for this effect may involve active chloride transport in the ascending limb of Henle, sodium absorption in the cortical collecting tubule or reduced glomerular filtration.<sup>138-141, 167, 168</sup> The involvement of prostaglandins in sodium excretion is most pronounced during diuretic administration. NSAIDs including aspirin and sulindac greatly impair the natriuretic response to furosemide, spironolactone and other diuretics.<sup>34, 169-171</sup> There is increasing evidence that prostaglandins are also involved in water excretion. A series of studies by Orloff and others suggested that prostaglandins produced in the collecting tubule may function as a negative feedback inhibitor on antidiuretic hormone-stimulated water reabsorption.<sup>172-174</sup> Although the interactions of antidiuretic hormone, cyclic AMP and prostaglandins are complex and not fully understood,<sup>175</sup> it is clear that NSAIDs contribute to water retention in humans and may contribute to dilutional hyponatremia. Alternatively, these drugs may be useful in reducing water loss in patients with central diabetes insipidus.<sup>138, 140</sup> NSAIDs also decrease water loss in patients with nephrogenic diabetes insipidus, indicating that the drugs also act independently of antidiuretic hormone, apparently by altering the medullary or glomerular blood flow.<sup>176</sup>

A series of animal studies by Morrison, Needleman and associates have shown that a ureteral obstructed kidney produces large amounts of thromboxane A<sub>2</sub>, which may increase renal vascular resistance and diminish renal blood flow.<sup>177, 178</sup> A similar situation occurs after partial obstruction of the renal vein.<sup>179</sup> In these models, fibroblast and mononuclear cell infiltration is the source of the augmented thromboxanes.<sup>180</sup> Renal thromboxane production is also increased in experimental glomerulonephritis.<sup>181</sup> The role of vasoconstrictor thromboxane in human renal function is less clear. Thromboxanes do not regulate renal blood flow in normal persons.<sup>182</sup> Urinary TxB<sub>2</sub> is greatly increased in patients with the hepatorenal syndrome,<sup>183</sup> but thromboxanes are probably not the cause of the renal vasoconstriction.<sup>184</sup> In patients with renal transplant rejection, urinary TxB<sub>2</sub> is also increased and serves as an early indication of rejection.<sup>185</sup> In this disorder it is likely that thromboxanes from infiltrating cells contribute to the rejection process. Platelet generation of thromboxanes or prostaglandins may also have deleterious long-term effects on the progression of membranoproliferative glomerulonephritis in humans.<sup>38</sup>

## Pulmonary Physiology and Diseases

In studies using pharmacologic amounts of eicosanoids in isolated tissues, animals and humans, possible functions of these chemicals have been suggested in regulating lung vascular and bronchial smooth muscle tone and in mucus secretion.<sup>186, 187</sup> Although there are species and preparation differences, pulmonary vascular constriction is caused by PGE<sub>2</sub> (opposite to its peripheral vasodilatory effect), PGD<sub>2</sub> and PGF<sub>2 $\alpha$</sub> .<sup>186-189</sup> PGE<sub>1</sub> is a vasodilator, and administering a PGE<sub>1</sub> analog to patients with chronic lung disease is effective in lowering pulmonary vascular resistance but does not immediately improve pulmonary function.<sup>190</sup> PGI<sub>2</sub> is also a potent

\*Reference numbers 143, 144, 151-153, 161-164.

vasodilator and weak bronchodilator, and infusions of PGI<sub>2</sub> have been administered for treating pulmonary hypertension.<sup>191</sup>

PGF<sub>2α</sub>, TxA<sub>2</sub> and PGD<sub>2</sub> are bronchoconstrictors, and the bronchoconstrictive effect of PGF<sub>2α</sub> aerosol is greatly potentiated in patients with asthma.<sup>187,192</sup> In contrast, PGE<sub>1</sub> and PGE<sub>2</sub> are bronchodilators and block the effects of PGF<sub>2α</sub>.<sup>193</sup> As described above, leukotrienes C<sub>4</sub>, D<sub>4</sub> and E<sub>4</sub> are extremely potent bronchoconstrictors, perhaps 1,000-fold more potent than histamine on bronchial tissue.<sup>194-196</sup> These leukotrienes also stimulate mucus secretion and increase vascular permeability.

These pharmacologic data are not synonymous with physiologic functions. However, there is strong evidence for eicosanoid activity in several types of pulmonary disease, particularly in patients with asthma.<sup>197</sup> SRS-A (now identified as peptide leukotrienes) has been implicated as a key mediator of allergic bronchospasm for decades.<sup>198</sup> These leukotrienes are synthesized in human lung tissue, especially from pulmonary mast cells.<sup>199,200</sup> Leukotriene generation is increased during an allergic challenge in patients with asthma in parallel with the onset of bronchoconstriction.<sup>201</sup> In contrast to their hypersensitive response to PGF<sub>2α</sub>, however, patients with asthma have the same degree of bronchoconstriction response to LTD<sub>4</sub> as do those who do not have asthma.<sup>202</sup> In guinea pigs, part of the leukotriene-induced bronchoconstriction is mediated by augmented TxA<sub>2</sub> release,<sup>203</sup> but in humans the leukotriene effect is independent of cyclooxygenase products.<sup>204</sup> About 15% of patients with asthma have an attack following ingestion of an NSAID, apparently related to cyclooxygenase inhibition<sup>205</sup> and possibly due to a shift in arachidonic acid metabolism from cyclooxygenase to lipoxygenase products. Corticosteroids are a well-established therapy for persistent asthma, and these drugs reduce lipoxygenase products, supporting the link between leukotrienes and asthma. Proof of this association and possible new therapeutic modalities must await development of specific leukotriene inhibitors.<sup>43</sup>

Eicosanoids have been implicated in several other types of acute pulmonary injury.<sup>206,207</sup> With massive pulmonary emboli, thromboxane release from activated platelets or lung parenchyma contributes to the hemodynamic alterations.<sup>208</sup> Assay of thromboxanes excreted in the urine may be a useful adjunct to diagnosing acute thromboembolic disease.<sup>209</sup> Thromboxanes and other cyclooxygenase products may also contribute to the pulmonary hypertension and respiratory distress that develops after trauma, fat emboli and possibly sepsis.<sup>206,207,210,211</sup> In studies of animals, cyclooxygenase and thromboxane inhibitors improve cardiopulmonary hemodynamics.<sup>210-213</sup> Infusions of PGI<sub>2</sub> or PGE<sub>1</sub> may also be useful by countering the effects of thromboxane, and clinical trials in humans are in progress.

Cyclooxygenase products are released into the circulation in large amounts during septic shock in animals and humans, and these products may also contribute to the systemic cardiovascular changes, including myocardial depression.<sup>213-217</sup> At least in some animal models of septic shock, pretreatment with thromboxane inhibitors improves survival.<sup>214,216,217</sup> The clinical usefulness of these agents for treating septic shock in humans has not been shown.

In inflammatory diseases of the lung, as with inflammation elsewhere, prostaglandins and leukotrienes are likely in-

volved. During inflammatory insults, such as bleomycin administration and exposure to asbestos fibers, an interaction between alveolar macrophages and fibroblasts involving prostaglandin E<sub>2</sub> and interleukin 1 may lead to collagen production and pulmonary fibrosis.<sup>218,219</sup>

### Gastrointestinal Physiology

Administering large amounts of prostaglandin E or F, such as for terminating a pregnancy, frequently causes diarrhea.<sup>220,221</sup> This phenomenon was initially attributed to direct stimulation of intestinal smooth muscle; it is now clear, however, that prostaglandins enhance intestinal fluid and electrolyte secretion and cause a secretory diarrhea.<sup>222,223</sup> Prostaglandins of the E and F type increase cyclic AMP in the small intestine and probably also in the colon, similar to the effects of cholera toxin.<sup>224,228</sup> Although still controversial, prostaglandins probably have only a minimal or secondary role in cholera toxin-induced diarrhea,<sup>229</sup> and prostaglandin inhibitors have not proved useful in treating human cholera infection. Endogenous prostaglandins may have a physiologic role in jejunal secretion in humans,<sup>230</sup> and augmented intestinal prostaglandin synthesis may contribute to the diarrhea associated with *Salmonella*, *Shigella* and *Escherichia coli* infections<sup>231</sup>; however, as in cholera infections, NSAIDs have not been clinically beneficial.

Prostaglandins have also been claimed to play a role in the diarrhea associated with a variety of noninfectious causes, including laxatives—such as bisacodyl, senna compounds, docusate sodium (dioctyl sodium sulfosuccinate)—thyrotoxicosis, hypergastrinemia, radiation-induced enteritis, irritable bowel syndrome and paraneoplastic syndromes.<sup>224-226</sup> It is difficult to interpret many of these early studies that suggested an association between prostaglandins and these clinical states. Some of the early prostaglandin assays of blood or intestinal contents were later shown to be problematic. The clinical response to NSAIDs often did not relate to increased basal prostaglandin levels. Studies done on animals were not always repeated in humans, and many publications were isolated case reports. In an excellent review, Metz, McRae and Robertson discuss the criteria to assess these publications.<sup>232</sup> Nevertheless, it is likely that overproduction of prostaglandins contributes to the diarrhea of a few patients with pancreatic tumors, medullary carcinoma of the thyroid and perhaps other conditions,<sup>232,233</sup> and a therapeutic trial of NSAIDs may be useful in patients occasionally seen who have diarrhea that is unresponsive to other modalities.

Prostaglandins are released at the sites of inflammation, including the colonic mucosa in cases of active inflammatory bowel disease.<sup>234-236</sup> These eicosanoids potentially may affect colonic motility, epithelial cell proliferation, anion secretion and inflammation.<sup>234-237</sup> However, administration of prostaglandin inhibitors does not appear to reduce the inflammation.<sup>238-240</sup> Conversely, administering prostaglandins does not appear to be effective in maintaining remission in patients with ulcerative colitis.<sup>241</sup> Lipoxygenase products are also increased in patients with colitis,<sup>242,243</sup> and these products may be more important than prostaglandins in contributing to inflammation and electrolyte secretion.<sup>244</sup> Recent studies by Stenson and Lobos have suggested that inhibition of leukotrienes is the mechanism of action of sulfasalazine and presumably corticosteroids in this disease.<sup>245</sup> The role of leuko-



trienes in inflammatory bowel disease will likely be further defined with the development of specific lipoxygenase inhibitors.

The most extensive research on prostaglandins and the gastrointestinal tract has focused on the gastric and duodenal mucosa. Early observations that exogenous prostaglandins inhibited gastric acid secretion in animals<sup>246</sup> and in humans<sup>247,248</sup> suggested that prostaglandins could be therapeutic agents for treating peptic ulcer disease. Synthetic prostaglandin analogs—especially 15,15- and 16,16-dimethyl PGE<sub>2</sub>—were more potent and longer acting than the natural prostaglandins. In studies using animals, exogenous prostaglandins were effective in healing duodenal ulcers and protecting against gastric mucosal injury induced by acid, alkali, base and a variety of other noxious stimuli.<sup>226–228,249,250</sup> These protective effects, however, were obtained with doses of prostaglandins less than 1/100th the dose needed to reduce acid secretion. The term “cytoprotection” is used to refer to this ability of exogenous prostaglandins to prevent damage to the gastric mucosa from noxious stimuli. Extensive research into the mechanism of cytoprotection has ensued. There are experimental data to support and to refute each of the proposed mechanisms: increased gastric mucosal blood flow, stimulation of cyclic AMP, prevention of gastric mucosal barrier disruption—that is, reducing hydrogen ion diffusion back into the mucosa—stimulation of mucus secretion, increase in alkaline secretion from nonparietal cells—that is, increasing the mucus-bicarbonate barrier—stimulation of cellular protein synthesis or transport or stabilization of lysosomes, maintenance of sulfhydryl compounds, increased cell proliferation and increase in mucosal surface hydrophobicity. It is clear from numerous clinical trials in humans that orally administered prostaglandins are equally effective as H<sub>2</sub> antagonists in healing duodenal ulcers, and prostaglandin analogs will likely be available for general clinical administration for this indication within a few years in this country. These compounds include 16,16-dimethyl PGE<sub>2</sub> (Upjohn), enprostil (Syntex), misoprostol (Searle), rioprostil (Ortho-Miles-Bayer) and trimoprostil (Hoffmann-La Roche).<sup>251</sup> These agents may also prove useful in treating gastric ulcer and hemorrhagic gastritis. The drugs are not without effects in other organs, and there is evidence that they may potentiate esophagitis associated with reduction in lower esophageal sphincter pressure.<sup>252,253</sup>

Endogenous production of prostaglandins from the gastric mucosa functions as a physiologic protective mechanism.<sup>226–228,249,250</sup> Current theory is that mild irritants, such as dilute ethanol, sodium chloride solutions and acid, stimulate prostaglandin production, which protects the gastric mucosa from subsequent necrotizing agents (more concentrated ethanol, for example). This phenomenon may also contribute to the therapeutic effects of sucralfate<sup>254</sup> and antacids.

Prostaglandins are likely involved in other aspects of gastrointestinal function. Gallbladder production of eicosanoids is increased in the inflamed tissues of cholecystitis and may contribute to decreased fluid absorption, gallbladder contraction and biliary pain.<sup>255</sup> Endogenous prostaglandins may also be involved in intestinal motility, although the clinical relevance to humans has not been shown. However, exogenous prostaglandins may be useful in stimulating motility in the treatment of postoperative ileus.<sup>256</sup>

## Metabolic Disorders

There is laboratory and hypothetical evidence implicating prostaglandins in many metabolic processes including hormone secretion, neurotransmission and mineral and electrolyte disorders. The topics of glucose regulation and hypercalcemia of malignancy have received the most attention.

The role of arachidonic acid metabolites in glucose regulation has been controversial.<sup>232,257–259</sup> Prostaglandin E is synthesized by the pancreatic islet, and infusions of prostaglandin E in humans generally inhibit insulin secretion and increase circulating glucose concentration. Although aspirin immediately enhances insulin release in persons with type II diabetes mellitus, available data fail to substantiate a clinically significant glucose-lowering effect in prolonged trials. Indomethacin has the opposite effect of other NSAIDs, apparently by a non-cyclooxygenase-mediated mechanism. Recent data suggest that a lipoxygenase product, probably 12-HETE, produced by the islet cells may have a key role in mediating insulin release.<sup>260</sup>

Prostaglandins were first implicated as mediators of hypercalcemia of malignancy in animal models.<sup>261</sup> Presumably a prostaglandin metabolite released into the circulation stimulates bone resorption, and indomethacin normalizes the hypercalcemia.<sup>261</sup> Recent findings have challenged this interpretation.<sup>262</sup> In humans, as many as 20% of cancer patients with hypercalcemia have no apparent bone metastasis, implicating humoral mechanisms for the hypercalcemia. An early study by Seyberth and associates implicated excessive circulating PGE, presumably released from the tumors, as the cause of hypercalcemia in many of these patients.<sup>263</sup> Administering indomethacin or aspirin reduced the serum calcium levels in the patients who had increased PGE metabolite levels. Other investigators have also shown increased prostaglandins and reduced serum calcium levels with indomethacin in a few patients<sup>264,265</sup>; it is not established, however, that circulating prostaglandins exist in high enough concentration to directly induce bone resorption. There is also evidence that the tumors may have products that induce the bone to synthesize eicosanoids, which then cause release of calcium.<sup>266</sup> At this time, the topic remains unclear, and circulating prostaglandins are probably not the explanation for the hypercalcemia of malignancy.<sup>267</sup>

Arachidonic acid metabolites have been implicated in other aspects of cancer, including immunosuppression, cancer cell growth and metastasis, and there is a potential use for NSAIDs in combination with chemotherapy. Several recent review articles discuss these topics.<sup>268–273</sup>

## Conclusions

Because of the recurring history of prostaglandin research in which many proposed physiologic functions could not be confirmed with subsequent better methods and study design, our concluding comment is to emphasize that readers carefully evaluate the assay techniques, the use of pharmacologic agonists and antagonists and the protocol design before accepting newly proposed physiologic functions for eicosanoids.

Research in arachidonic acid metabolites has increased greatly over the past two decades. The discovery of leukotrienes and related compounds will likely lead to clinical benefit in patients with asthma and inflammatory conditions. In

1985, prostaglandins have definitely become part of the clinical practice of medicine, particularly in the fields of reproduction, gastroenterology, nephrology and rheumatology.

# REFERENCES

1. Kurzrok R, Lieb CC: Biochemical studies of human semen—II. The action of semen on the human uterus. *Proc Soc Exp Biol Med* 1930; 28:268-272
2. Goldblatt MW: Properties of human seminal plasma. *J Physiol* 1935; 84:208-218
3. Von Euler US: Zur kenntnis der pharmakologischen Wirkungen von natirsekreten und Extrakten männlicher accessorischer Geschlechtsdrüsen. *Arch Exp Path Pharmacol* 1934; 175:78-84
4. Von Euler US: Über die spezifische blutdrucksenkende Substanz des menschlichen Prostata- und Samenblanzsekretes. *Klin Wochenschr* 1935; 14:1182-1183
5. Feldberg W, Kellaway CH: Liberation of histamine and formation of lysocithin-like substances by cobra venom. *J Physiol* 1938; 94:187-226
6. Kellaway CH, Trethowie EF: The liberation of a slow-reacting smooth muscle stimulating substance in anaphylaxis. *Q J Exp Physiol Cogn Med Sci* 1940; 30:121-145
7. Bergström S, Danielsson H, Samuelsson B: The enzymatic formation of prostaglandin E<sub>2</sub> from arachidonic acid. *Biochem Biophys Acta* 1964; 90:207-210
8. Van Dorp DA, Beerthuis RK, Nugteren DH, et al: The biosynthesis of prostaglandins. *Biochim Biophys Acta* 1964; 90:204-207
9. Vane JR: Inhibition of prostaglandin synthesis as a mechanism of action for aspirin-like drugs. *Nature (New Biol)* 1971; 231:232-235
10. Hamberg M, Svensson J, Samuelsson B: Thromboxanes: A new group of biologically active compounds derived from prostaglandin endoperoxides. *Proc Natl Acad Sci USA* 1975; 72:2994-2998
11. Moncada S, Gryglewski R, Bunting S, et al: An enzyme isolated from arteries transforms prostaglandin endoperoxides to an unstable substance that inhibits platelet aggregation. *Nature* 1976; 263:663-665
12. Hammarström S: Leukotrienes. *Annu Rev Biochem* 1983; 52:355-377
13. Borgeat P, Samuelsson B: Arachidonic acid metabolism in polymorphonuclear leukocytes: Effects of ionophore A23187. *Proc Natl Acad Sci USA* 1979; 76:2148-2152
14. Murphy RC, Hammarström S, Samuelsson B: Leukotriene C: A slow-reacting substance from murine mastocytoma cells. *Proc Natl Acad Sci USA* 1979; 76:4275-4279
15. Dyerberg J, Bang HO, Stoffersen E, et al: Eicosapentaenoic acid and prevention of thrombosis and atherosclerosis? *Lancet* 1978; 2:117-119
16. Fridovich I: The biology of oxygen radicals. *Science* 1978; 201:875-880
17. Siegel MI, McConnell RT, Cuatrecasas P: Aspirin-like drugs interfere with arachidonate metabolism by inhibition of the 12-hydroperoxy-5,8,10,14-eicosatetraenoic acid peroxidase activity of the lipoxygenase pathway. *Proc Natl Acad Sci USA* 1979; 76:3774-3778
18. Samuelsson B, Granström E, Green K, et al: Prostaglandins. *Annu Rev Biochem* 1975; 44:669-695
19. Samuelsson B, Goldyne M, Granström E, et al: Prostaglandins and thromboxanes. *Annu Rev Biochem* 1978; 47:997-1029
20. Lands WEM: The biosynthesis and metabolism of prostaglandins. *Annu Rev Physiol* 1979; 41:633-652
21. Samuelsson B: Leukotrienes: Mediators of immediate hypersensitivity reactions and inflammation. *Science* 1983; 220:568-575
22. Lewis RA, Austen KF: The biologically active leukotrienes: Biosynthesis, metabolism, receptors, functions, and pharmacology. *J Clin Invest* 1984; 73:889-897
23. Samuelsson F: From studies of biochemical mechanism to novel biological mediators: Prostaglandin endoperoxides, thromboxanes, and leukotrienes—Nobel Lecture, 8 December 1982. *Biosci Rep* 1983; 3:791-813
24. Samuelsson B, Hammarström S: Leukotrienes: A novel group of biologically active compounds. *Vitam Horm* 1982; 39:1-30
25. Serhan CN, Hamberg M, Samuelsson B: Lipoxins: Novel series of biologically active compounds formed from arachidonic acid in human leukocytes. *Proc Natl Acad Sci USA* 1984; 81:5335-5339
26. Simon LS, Mills JA: Drug therapy: Nonsteroidal anti-inflammatory drugs. *N Engl J Med* 1980; 302:1179-1185, 1237-1243
27. Huskisson EC (Ed): *Anti-Rheumatic Drugs*. New York, Praeger Scientific, 1983, pp 1-749
28. Weissmann G: Prostaglandins in acute inflammation. *In Current Concepts*. Kalamazoo, Mich, A Scope Publication of the Upjohn Company, 1980, pp 1-32
29. Metz SA: Anti-inflammatory agents as inhibitors of prostaglandin synthesis in man. *Med Clin North Am* 1981; 65:713-757
30. Siegel MI, McConnell RT, Porter NA, et al: Aspirin-like drugs inhibit arachidonic acid metabolism via lipoxygenase and cyclo-oxygenase in rat neutrophils from carrageenan pleural exudates. *Biochem Biophys Res Commun* 1980; 92:688-695
31. Robinson DR: Prostaglandins and anti-inflammatory drugs. *DM* 1983; 30:2-46
32. Flower RJ: Drugs which inhibit prostaglandin biosynthesis. *Pharmacol Rev* 1974; 26:33-67
33. Ciabattini G, Cinotti GA, Pierucci A, et al: Effects of sulindac and ibuprofen in patients with chronic glomerular disease. *N Engl J Med* 1984; 310:279-283
34. Daskalopoulos G, Kronborg I, Katkov W, et al: Sulindac and indomethacin suppress the diuretic action of furosemide in patients with cirrhosis and ascites: Evidence that sulindac affects renal prostaglandins. *Am J Kidney Dis*, in press
35. Whittle BJR, Higgs GA, Eakins KE, et al: Selective inhibition of prostaglandin production in inflammatory exudates and gastric mucosa. *Nature* 1980; 284:271-273
36. Needleman P, Raz A, Ferrendelli JA, et al: Application of imidazole as a selective inhibitor of thromboxane synthetase in human platelets. *Proc Natl Acad Sci USA* 1977; 74:1716-1720
37. Tyler HM, Saxton CAPD, Parry MJ: Administration to man of UK-37,248-01, a selective inhibitor of thromboxane synthetase. *Lancet* 1981; 1:629-632
38. Donadio JV Jr, Anderson CF, Mitchell JC III, et al: Membranoproliferative glomerulonephritis: A prospective clinical trial of platelet-inhibitor therapy. *N Engl J Med* 1984; 310:1421-1426
39. Hong SL, Levine L: Inhibition of arachidonic acid release from cells as the biochemical action of anti-inflammatory corticosteroids. *Proc Natl Acad Sci USA* 1976; 73:1730-1734
40. Blackwell GJ, Carnuccio R, DiRosa M, et al: Macrocortin: A polypeptide causing the anti-phospholipase effect of glucocorticoids. *Nature* 1980; 287:147-149
41. Hirata F: The regulation of lipomodulin, a phospholipase inhibitory protein, in rabbit neutrophils by phosphorylation. *J Biol Chem* 1981; 256:7730-7733
42. DiRosa M, Flower RJ, Hirata F, et al: Nomenclature announcement: Anti-phospholipase proteins. *Prostaglandins* 1984; 28:441
43. Bach MK: Prospects for the inhibition of leukotriene synthesis. *Biochem Pharmacol* 1984; 33:515-521
44. Granström E, Kindahl H: Radioimmunoassay of prostaglandins and thromboxanes. *In Frolich JC (Ed): Advances in Prostaglandin and Thromboxane Research*, Vol 5. New York, Raven Press, 1978, pp 119-210
45. FitzGerald GA, Pedersen AK, Patrono C: Analysis of prostacyclin and thromboxane biosynthesis in cardiovascular disease. *Circulation* 1983; 67:1174-1177
46. FitzGerald GA, Oates JA, Hawiger J, et al: Endogenous biosynthesis of prostacyclin and thromboxane and platelet function during chronic aspirin administration in man. *J Clin Invest* 1983; 71:676-688
47. Granström E, Kindahl H, Swann ML: Profiles of prostaglandin metabolites in the human circulation—Identification of late-appearing long-lived products. *Biochim Biophys Acta* 1982; 713:46-60
48. Lee JB, Crowshaw K, Takman BH, et al: The identification of prostaglandins E<sub>2</sub>, F<sub>2α</sub>, and A<sub>2</sub> from rabbit kidney medulla. *Biochem J* 1967; 105:1251-1260
49. Zusman RM, Spector D, Caldwell BV, et al: The effect of chronic sodium loading and sodium restriction on plasma prostaglandin A, E, and F concentrations in normal humans. *J Clin Invest* 1973; 52:1093-1098
50. Lee JB: Renal homeostasis and the hypertensive state: A unifying hypothesis. *In Ramwell PW (Ed): The Prostaglandins*, Vol 1. New York, Plenum Press, 1973, pp 133-187
51. Frolich JC, Sweetman BJ, Carr K, et al: Assessment of the levels of PGA<sub>2</sub> in human plasma by gas chromatography-mass spectrometry. *Prostaglandins* 1975; 10:185-195
52. FitzGerald GA, Brash AR, Falardeau P, et al: Estimated rate of prostacyclin secretion into the circulation of normal man. *J Clin Invest* 1981; 86:1271-1276
53. Martin K, Zipser RD, Horton R: The effect of prostaglandin inhibition on the hypertensive action of sodium retaining steroids. *Hypertension* 1981; 3:622-628
54. Gerber JG: Indomethacin-induced rises in blood pressure. *Ann Intern Med* 1983; 99:555-558
55. Dzau VJ, Packer M, Lilly LS, et al: Prostaglandins in severe congestive heart failure: Relation to activation of the renin angiotensin system and hyponatremia. *N Engl J Med* 1984; 310:347-352
56. Packham MA: Mode of action of acetylsalicylic acid. *In Barnett HJM, Hirsh J, Mustard JF (Eds): Acetylsalicylic Acid: New Uses for an Old Drug*. New York, Raven Press, 1982, pp 63-86
57. Whittle BJR, Moncada S: Pharmacology of prostacyclin and thromboxanes. *Br Med Bull* 1983; 39:232-238
58. Silver MJ: Mechanisms of hemostasis and therapy of thrombosis: New concepts based on the metabolism of arachidonic acid by platelets and endothelial cells. *Adv Pharmacol Chemother* 1981; 18:1-47
59. DiMinno G, DeGaetano G, Garattini S: Dipyridamole and platelet function. *Lancet* 1978; 2:1258-1259
60. Harlan JM, Harker LA: Hemostasis, thrombosis, and thromboembolic disorders: The role of arachidonic acid metabolites in platelet-vessel wall interactions. *Med Clin North Am* 1981; 65:855-880
61. Cannon PJ: Eicosanoids and the blood vessel wall. *Circulation* 1984; 70:523-528
62. Gryglewski RJ: Prostacyclin, prostaglandins, thromboxanes, and platelet function. *In Lee JL (Ed): Prostaglandins*. New York, Elsevier, 1981, pp 303-350
63. Stuart MJ, Gerrard JM, White JG: Effect of cholesterol on production of thromboxane B<sub>2</sub> by platelets in vitro. *N Engl J Med* 1980; 302:6-10
64. Halushka PV, Rogers RC, Loadholt CB, et al: Increased platelet thromboxane synthesis in diabetes mellitus. *J Lab Clin Med* 1981; 97:87-96
65. Pomerantz KB, Tall AR, Feinmark SJ, et al: Stimulation of vascular smooth muscle cell prostacyclin synthesis by high density lipoproteins. *Circ Res* 1984; 54:554-565
66. Nadler JL, Velasco JS, Horton R: Cigarette smoking inhibits prostacyclin formation. *Lancet* 1983; 1:1248-1250
67. FitzGerald GA, Smith B, Pedersen AK, et al: Increased prostacyclin biosynthesis in patients with severe atherosclerosis and platelet activation. *N Engl J Med* 1984; 310:1065-1068
68. Eichner ER: Platelets, carotids, and coronaries: Critique on antithrombotic role of antiplatelet agents, exercise, and certain diets. *Am J Med* 1984; 77:513-523
69. Marcus AJ: Aspirin as an antithrombotic medication. *N Engl J Med* 1983; 309:1515-1516
70. Marcus AJ: Arachidonate metabolism in vascular disorders. *J Clin Invest* 1983; 72:1521-1525
71. Chesebro JH, Fuster V, Elveback LR, et al: Effect of dipyridamole and aspirin

- in late vein-graft patency after coronary bypass operations. *N Engl J Med* 1984; 310:209-214
72. Pitt B, Shea M, Romson J, et al: Prostaglandins and prostaglandin inhibitors in ischemic heart disease. *Ann Intern Med* 1983; 99:83-92
73. Vane JR: Clinical potential of prostacyclin. *In* Samuelsson B, Paoletti R, Ramwell P (Eds): *Advances in Prostaglandin, Thromboxane, and Leukotriene Research*, Vol 11. New York, Raven Press, 1983, pp 449-456
74. Bunting S, O'Grady J, Fabiana JN, et al: Cardiopulmonary bypass in man: Effects of prostacyclin. *In* Lewis PJ, O'Grady JM (Eds): *Clinical Pharmacology of Prostacyclin*. New York, Raven Press, 1981, pp 181-193
75. Turney JH, Williams LC, Fewell MR, et al: Platelet protection and heparin sparing with prostacyclin during regular dialysis therapy. *Lancet* 1980; 2:219-222
76. Gimson AE, Langley PG, Huges RD, et al: Prostacyclin to prevent platelet activation during charcoal haemoperfusion in fulminant hepatic failure. *Lancet* 1980; 1:173-175
77. Walker ID, Davidson JF, Faichney A, et al: A double-blind study of prostacyclin in cardiopulmonary bypass surgery. *Br J Haematol* 1981; 49:415-423
78. Szczeklik A, Nizankowski R, Skawinski S, et al: Successful therapy of advanced arteriosclerosis obliterans with prostacyclin. *Lancet* 1979; 1:1111-1114
79. Gryglewski RJ, Nowak S, Kostka-Trabka E, et al: Treatment of ischaemic stroke with prostacyclin. *Stroke* 1983; 14:197-202
80. Carlson LA, Olsson AG: PGE<sub>1</sub> in ischaemic peripheral vascular disease. *In* Karim SMM (Ed): *Practical Applications of Prostaglandins and Their Synthesis Inhibitors*. Baltimore, University Park Press, 1979, pp 39-51
81. Dowd PM, Martin MF, Cooke ED, et al: Treatment of Raynaud's phenomenon by intravenous infusion of prostacyclin (PGI<sub>2</sub>). *Br J Dermatol* 1982; 106:81-89
82. Belch JJ, McKay A, McArdle B, et al: Epoprostenol (prostacyclin) and severe arterial disease—A double-blind trial. *Lancet* 1983; 1:315-317
83. Belch JJ, Newman P, Drury JK, et al: Intermittent epoprostenol (prostacyclin) infusion in patients with Raynaud's syndrome—A double-blind controlled trial. *Lancet* 1983; 1:313-315
84. Hossman V, Heinen A, Auel H, et al: A randomized placebo controlled trial of prostacyclin in peripheral arterial disease of the lower extremities. *Thromb Res* 1981; 22:481-490
85. Goldman DW, Goetzl EJ: Mediation and modulation of immediate hypersensitivity and inflammation byproducts of the oxygenation of arachidonic acid. *In* Ward PA (Ed): *Immunology of Inflammation: Handbook of Inflammation*, Vol 4. Amsterdam, Elsevier, 1983, pp 163-187
86. Zurier RB: Prostaglandins and inflammation. *In* Lee JB (Ed): *Prostaglandins*. New York, Elsevier, 1982, pp 91-112
87. Stenson WJ, Parker CW: Prostaglandins and the immune response. *In* Lee JB (Ed): *Prostaglandins*. New York, Elsevier, 1982, pp 39-89
88. Goodwin JS, Ceuppens J: Regulation of the immune response by prostaglandins. *J Clin Immunol* 1983; 3:295-315
89. Simmons PM, Salmon JA, Moncada S: The release of leukotriene B<sub>4</sub> during experimental inflammation. *Biochem Pharmacol* 1983; 32:1353-1359
90. Robinson DR, Dayer J-M, Krane SM: Prostaglandins and their regulation in rheumatoid inflammation. *Ann NY Acad Sci* 1979; 332:279-294
91. Korn JH, Halushka PV, LeRoy EC: Mononuclear cell modulation of connective tissue function—Suppression of fibroblast growth by stimulation of endogenous prostaglandin production. *J Clin Invest* 1980; 65:543-554
92. Kuehl FA Jr, Humes JL, Egan RW, et al: Role of prostaglandin endoperoxide PGG<sub>2</sub> in inflammatory processes. *Nature* 1977; 265:170-173
93. Baracos V, Rodemann HP, Dinarello CA, et al: Stimulation of muscle protein degradation and prostaglandin E<sub>2</sub> release by leukocytic pyrogen (interleukin-1)—A mechanism for the increased degradation of muscle protein during fever. *N Engl J Med* 1983; 308:553-558
94. Beisel WR: Mediators of fever and muscle proteolysis (Editorial). *N Engl J Med* 1983; 308:586-587
95. Gimbrone MA Jr, Brock AF, Schafer AI: Leukotriene B<sub>4</sub> stimulates polymorphonuclear leukocyte adhesion to cultured vascular endothelial cells. *J Clin Invest* 1984; 74:1552-1555
96. Brain S, Camp R, Dowd P, et al: The release of leukotriene B<sub>4</sub>-like material in biologically active amounts from the lesional skin of patients with psoriasis. *J Invest Dermatol* 1984; 83:70-73
97. Hammarström S, Hamberg M, Samuelsson B, et al: Increased concentration of free arachidonic acid, prostaglandins E<sub>2</sub> and F<sub>2α</sub> and of 12L-hydroxy-5,8,10,14-eicosatetraenoic acid (HETE) in epidermis of psoriasis: Evidence for perturbed regulation of arachidonic acid levels in psoriasis. *Proc Natl Acad Sci USA* 1975; 72:5130-5134
98. Sharon P, Stenson WF: Enhanced synthesis of leukotriene B<sub>4</sub> by colonic mucosa in inflammatory bowel disease. *Gastroenterology* 1984; 86:453-460
99. Klickstein LB, Shapleigh C, Goetzl EJ: Lipooxygenation of arachidonic acid as a source of polymorphonuclear leukocyte chemotactic factor in synovial fluid and in tissue in rheumatoid arthritis and spondyloarthritis. *J Clin Invest* 1980; 66:1166-1170
100. Roberts LJ II, Sweetman BJ, Lewis RA, et al: Increased production of prostaglandin D<sub>2</sub> in patients with systemic mastocytosis. *N Engl J Med* 1980; 303:1400-1404
101. Gordon D, Bray MA, Morley J: Control of lymphokine secretion by prostaglandins. *Nature (Lond)* 1976; 262:401-402
102. Goodwin JS, Messner RP, Bankhurst AD, et al: Prostaglandin producing suppressor cells in Hodgkin's disease. *N Engl J Med* 1977; 297:963-968
103. Goodwin JS: Immunologic effects of nonsteroidal anti-inflammatory drugs. *Am J Med* 1984; 77(4B):7-15
104. Karim SMM: *Practical Applications of Prostaglandins*. Baltimore, University Park Press, 1979
105. Gail J: The use of PGs in human reproduction. *Popul Rep [G]* 1980; 8:77-118
106. Patrick JE, Challis JRG: The role of prostaglandins and their inhibitors in reproduction. *In* Barnett HJM, Hirsh J, Mustard JF (Eds): *Acetylsalicylic Acid: New Uses for an Old Drug*. New York, Raven Press, 1982, pp 123-136
107. Caldwell BV, Behrman HR: Prostaglandins in reproductive processes. *Med Clin North Am* 1981; 65:927-936
108. Samuelsson B: Isolation and identification of prostaglandins from human seminal plasma. *J Biol Chem* 1963; 238:3229-3234
109. Cooper J, Kelly RW: The measurement of E and 19-hydroxy E prostaglandins in human seminal fluid. *Prostaglandins* 1975; 10:507-514
110. Bygdeman M, Fredricsson B, Svanborg K, et al: The relation between fertility and prostaglandin content of seminal fluid in man. *Fertil Steril* 1970; 21:622-630
111. Armstrong DT, Grinwich DL, Moon YS, et al: Inhibition of ovulation in rabbits by intrafollicular injection of indomethacin and prostaglandin F antiserum. *Life Sci* 1974; 14:129-140
112. O'Grady JP, Caldwell BV, Auletta EJ, et al: The effects of an inhibitor of prostaglandin synthesis (indomethacin) on ovulation, pregnancy and pseudopregnancy in the rabbit. *Prostaglandins* 1972; 1:97-106
113. Denefors B, Hamberger L, Hillensjö T, et al: Aspects concerning the role of prostaglandins for ovarian function. *Acta Obstet Gynecol Scand* 1983; 113(suppl):31-41
114. Dawson W, Lewis RL, McMahon RE, et al: Potent luteolytic agents related to prostaglandin F<sub>2α</sub>. *Nature* 1974; 250:330-331
115. Horton EW, Poyster NL: Uterine luteolytic hormone: A physiological role for prostaglandin F<sub>2α</sub>. *Physiol Rev* 1976; 56:595-651
116. Cooper MJ, Hammond D, Schultz RH: Veterinary uses of prostaglandins. *In* Karim SMM (Ed): *Practical Applications of Prostaglandins and Their Synthesis Inhibitors*. Baltimore, University Park Press, 1979, pp 189-216
117. Chan WY: Prostaglandins and nonsteroidal antiinflammatory drugs in dysmenorrhea. *Annu Rev Pharmacol Toxicol* 1983; 23:131-149
118. Wqvist N: Prostaglandins and their synthesis inhibitors in primary dysmenorrhea. *In* Karim SMM (Ed): *Practical Applications of Prostaglandins and Their Synthesis Inhibitors*. Baltimore, University Park Press, 1979, pp 217-265
119. Pulkkinen MO: Prostaglandins and the non-pregnant uterus—The pathophysiology of primary dysmenorrhea. *Acta Obstet Gynecol Scand* 1983; 113(suppl):63-67
120. Owen PR: Prostaglandin synthetase inhibitors in the treatment of primary dysmenorrhea: Outcome trials reviewed. *Am J Obstet Gynecol* 1984; 148:96-103
121. Anderson ABM, Haynes PJ, Guillebaud J, et al: Reduction of menstrual blood loss by prostaglandin synthetase inhibitors. *Lancet* 1976; 1:774-776
122. Topozada M: Prostaglandins and their synthesis inhibitors in dysfunctional uterine bleeding. *In* Karim SMM (Ed): *Practical Applications of Prostaglandins and Their Synthesis Inhibitors*. Baltimore, University Park Press, 1979, pp 237-265
123. Christensen NJ, Green K: Endogenous prostaglandin synthesis and abortion: A mini review. *Acta Obstet Gynecol Scand* 1983; 113(suppl):109-111
124. Brenner WE, Berger GS: Pharmacologic methods of inducing mid-trimester abortion: Risks and benefits. *In* Sciarra JJ, Zatzuchni GI, Speidel JJ (Eds): *Risks, Benefits, and Controversies in Fertility Control*. New York, Harper & Row, 1978, pp 292-321
125. Kajanoja P: Induction of abortion by prostaglandins in the second trimester of pregnancy: A review. *Acta Obstet Gynecol Scand* 1983; 113(suppl):145-151
126. Karim SMM: Termination of second trimester pregnancy with prostaglandins. *Practical Applications of Prostaglandins and Their Synthesis Inhibitors*. Baltimore, University Park Press, 1979, pp 375-409
127. Karim SMM, Ng SC, Ratnam SS: Termination of abnormal intrauterine pregnancy with prostaglandins. *In* Karim SMM (Ed): *Practical Applications of Prostaglandins and Their Synthesis Inhibitors*. Baltimore, University Park Press, 1979, pp 319-374
128. Karim SMM, Trussell RR, Patel RC, et al: Response of pregnant human uterus to prostaglandin F<sub>2α</sub>: Induction of labor. *Br Med J* 1968; 4:621-623
129. Lange AP, Westergaard JG, Secher NJ, et al: Labor induction with prostaglandins. *Acta Obstet Gynecol Scand* 1983; 113(suppl):177-185
130. Levin DL: Effects of inhibition of prostaglandin synthesis on fetal development, oxygenation and the fetal circulation. *Semin Perinatol* 1980; 4:35-44
131. Ellwood DA, Mitchell MD, Andersson A, et al: Oestrogens, prostaglandins and cervical ripening. *Lancet* 1979; 1:376-377
132. Calder AA: Prostaglandins for pre-induction cervical ripening. *In* Karim SMM (Ed): *Practical Applications of Prostaglandins and Their Synthesis Inhibitors*. Baltimore, University Park Press, 1979, pp 301-318
133. Heymann MA, Rudolph AM, Silverman NH: Closure of the ductus arteriosus in premature infants by inhibition of prostaglandin synthesis. *N Engl J Med* 1976; 295:530-533
134. Friedman WF, Hirschklau MK, Printz MP, et al: Pharmacological closure of the patent ductus arteriosus in premature infants. *N Engl J Med* 1976; 295:526-529
135. Coceani F, Olley PM: Action of prostaglandin synthetase inhibitors on the ductus arteriosus: Experimental and clinical aspects. *In* Barnett HJM, Hirsh J, Mustard JF (Eds): *Acetylsalicylic Acid: New Uses for an Old Drug*. New York, Raven Press, 1982, pp 109-122
136. Heymann MA: Pharmacologic use of prostaglandin E<sub>1</sub> in infants with congenital heart disease. *Am Heart J* 1981; 101:837-843
137. Printz MP, Skidgel RA, Friedman WF: Studies of pulmonary prostaglandin biosynthetic and catabolic enzymes as factors in ductus arteriosus patency and closure—Evidence for a shift in products with gestational age. *Pediatr Res* 1984; 18:19-24
138. Dunn MJ, Patrono C, Cinotti GA (Eds): *Prostaglandins and the Kidney*. New York, Plenum Medical Book, 1983
139. Dunn MJ: Renal prostaglandins. *Renal Endocrinology*. Baltimore, Williams & Wilkins, 1983
140. Horton R, Zipser R, Fichman M: Prostaglandins, renal function and vascular regulation. *Med Clin North Am* 1981; 65:891-914

141. Ferris TF: Prostaglandins and the kidney. *Am J Nephrol* 1983; 3:139-144
142. Levenson DJ, Simmons CE, Brenner BM: Arachidonic acid metabolism, prostaglandins and the kidney. *Am J Med* 1982; 72:354-374
143. Clive DM, Stoff JS: Renal syndromes associated with nonsteroidal anti-inflammatory drugs. *N Engl J Med* 1984; 310:563-572
144. Garella S, Matarese RA: Renal effects of prostaglandins and clinical adverse effects of nonsteroidal anti-inflammatory agents. *Medicine (Baltimore)* 1984; 63:165-181
145. Verberkmoes R, van Damme B, Clement J, et al: Bartter's syndrome with hyperplasia of the renomedullary interstitial cells—Successful treatment with indomethacin. *Kidney Int* 1976; 9:302-307
146. Fichman MP, Telfer N, Zia P, et al: Role of prostaglandins in the pathogenesis of Bartter's syndrome. *Am J Med* 1976; 60:785-797
147. Zipser RD, Rude RK, Zia PK, et al: Regulation of urinary prostaglandins in Bartter's syndrome. *Am J Med* 1979; 67:263-267
148. Gill JR, Bartter FC: Evidence for a prostaglandin-independent defect in chloride reabsorption in the loop of Henle as a proximal cause of Bartter's syndrome. *Am J Med* 1978; 65:766-772
149. Rumpf KW, Frenzel S, Lowitz HD, et al: The effect of indomethacin on plasma renin activity in man under normal conditions and after stimulation of the renin-angiotensin system. *Prostaglandins* 1975; 10:641-648
150. Veldhuis JD, Bardin CW, Demers LM: Metabolic mimicry of Bartter's syndrome by covert vomiting: Utility of urinary chloride determinations. *Am J Med* 1979; 66:361-363
151. Boyer TD, Zia P, Reynolds TB: Effect of indomethacin and prostaglandin A<sub>1</sub> on renal function and plasma renin activity in alcoholic liver disease. *Gastroenterology* 1979; 77:215-222
152. Zipser RD, Hoefs J, Speckart P, et al: Prostaglandins: Modulators of renal function and pressor resistance in chronic liver disease. *J Clin Endocrinol Metab* 1979; 48:895-900
153. Gentilini P, Laffi G, Buzzelli J: Functional renal alterations in chronic liver disease. *Digestion* 1980; 20:73-78
154. Gerber JG, Olson RD, Nies AS: Interrelationship between prostaglandins and renin release. *Kidney Int* 1981; 19:816-821
155. Horton R: Prostaglandins and the renin-angiotensin system. *Miner Electrolyte Metab* 1981; 6:60-67
156. Henrich WL: Role of prostaglandins in renin secretion. *Kidney Int* 1981; 19:822-830
157. Freeman RH, Davis JO, Villarreal D: Role of renal prostaglandins in the control of renin release. *Circ Res* 1984; 54:1-9
158. Patrono C, Pugliese F, Ciabattoni G, et al: Evidence for a direct stimulatory effect of prostacyclin on renin release in man. *J Clin Invest* 1982; 69:231-239
159. Speckart P, Zia P, Zipser R, et al: The effect of sodium restriction and prostaglandin inhibition on the renin-angiotensin system in man. *J Clin Endocrinol Metab* 1977; 44:832-837
160. Tan SY, Shapiro R, Franco R, et al: Indomethacin-induced prostaglandin inhibition with hyperkalemia. *Ann Intern Med* 1979; 90:783-785
161. Tan SY, Shapiro R, Kish MA: Reversible acute renal failure induced by indomethacin. *JAMA* 1979; 241:2732-2733
162. Donker AJM, Arisz L, Brentjens JRH, et al: The effect of indomethacin on kidney function and plasma renin activity in man. *Nephron* 1976; 17:788-796
163. Kimberly RP, Plotz PH: Aspirin-induced depression of renal function. *N Engl J Med* 1977; 296:418-428
164. Stillman MT, Napier J, Blackshear JL: Adverse effects of nonsteroidal anti-inflammatory drugs on the kidney. *Med Clin North Am* 1984; 68:371-385
165. Donker ABJ: The effect of indomethacin on renal function and glomerular protein loss. *In* Dunn MJ, Patrono C, Cinotti GA (Eds): *Prostaglandins and the Kidney*. New York, Plenum Medical Book, 1983, pp 251-262
166. Zipser RD, Zia P, Stone RA, et al: The prostaglandin and kallikrein-kinin systems in mineralocorticoid escape. *J Clin Endocrinol Metab* 1978; 47:996-1001
167. Stokes JB, Kokko JP: Inhibition of sodium transport by prostaglandin E<sub>2</sub> across the isolated perfused rabbit collecting tubule. *J Clin Invest* 1977; 59:1099-1104
168. Higashihara E, Stokes JB, Kokko JP, et al: Cortical and papillary micropuncture examination of chloride in segments of rat kidney during inhibition of prostaglandin production. *J Clin Invest* 1979; 64:1277-1287
169. Gerber JG: Role of prostaglandins in the hemodynamic and tubular effects of furosemide. *Fed Proc* 1983; 42:1707-1710
170. Patak RV, Mookerjee BK, Bentzel CJ, et al: Antagonism of the effects of furosemide by indomethacin in normal and hypertensive man. *Prostaglandins* 1975; 10:649-659
171. Mirouze D, Zipser RD, Reynolds T: Inhibition of the action of diuretics by non-steroidal anti-inflammatory drugs in patients with chronic liver disease and ascites. *Hepatology* 1983; 3:50-55
172. Orloff J, Handler JS, Bergstrom S: Effect of prostaglandin (PGE<sub>1</sub>) on the permeability response of the toad bladder to vasopressin, theophylline and adenosine 3':5'-monophosphate. *Nature (Lond)* 1965; 205:397-398
173. Zusman RM, Keiser HR, Handler JS: Vasopressin-stimulated prostaglandin E biosynthesis in the toad urinary bladder: Effect on water flow. *J Clin Invest* 1977; 60:1339-1347
174. Zusman RM: Prostaglandins, vasopressin, and renal water reabsorption. *Med Clin North Am* 1981; 65:915-925
175. Garcia-Perez A, Smith WL: Apical-basolateral membrane asymmetry in canine cortical tubule cells—Bradykinin, arginine vasopressin, prostaglandin E<sub>2</sub> interrelationships. *J Clin Invest* 1984; 74:63-74
176. Usberti M, Dechaux M, Guillot M, et al: Renal prostaglandin E<sub>2</sub> in nephrogenic diabetes insipidus: Effects of inhibition of prostaglandin synthesis by indomethacin. *J Pediatr* 1980; 97:476-478
177. Morrison AR, Nishikawa K, Needleman P: Unmasking of thromboxane A<sub>2</sub> synthesis by ureteral obstruction in the rabbit kidney. *Nature* 1977; 267:259-260
178. Currie M, Kawasaki A, Jonas P, et al: The mechanism and site of the enhanced arachidonate metabolism in ureter obstruction. *In* Dunn MJ, Patrono C, Cinotti GA (Eds): *Prostaglandins and the Kidney—Biochemistry, Physiology, Pharmacology and Clinical Applications*. New York, Plenum Press, 1983, pp 299-308
179. Zipser RD, Myers S, Needleman P: Exaggerated prostaglandin and thromboxane synthesis in the rabbit with renal vein constriction. *Circ Res* 1980; 47:231-237
180. Okegawa T, Jonas PE, DeSchryver K, et al: Metabolic and cellular alterations underlying the exaggerated renal prostaglandin and thromboxane synthesis in ureter obstruction in rabbits: Inflammatory response involving fibroblasts and mononuclear cells. *J Clin Invest* 1983; 71:81-90
181. Lianos EA, Andres GA, Dunn MJ: Glomerular prostaglandin and thromboxane synthesis in rat nephrotic serum nephritis: Effects on renal hemodynamics. *J Clin Invest* 1983; 72:1439-1448
182. Zipser RD: The effects of selective inhibition of thromboxane synthesis on renal function in man. *Am J Physiol* 1985, in press
183. Zipser RD, Radvan G, Kronborg I, et al: Urinary thromboxane B<sub>2</sub> and prostaglandin E<sub>2</sub> in the hepatorenal syndrome: Evidence for increased vasoconstrictor and decreased vasodilator factors. *Gastroenterology* 1982; 84:697-703
184. Zipser RD, Kronborg I, Radvan G, et al: Therapeutic trial of thromboxane synthesis inhibition in the hepatorenal syndrome. *Gastroenterology* 1984; 87:1228-1232
185. Foegh ML, Winchester JF, Zmudka M, et al: Urine i-TxB<sub>2</sub> in renal allograft rejection. *Lancet* 1981; 2:431-434
186. Mathe AA, Hedqvist P, Strandberg K, et al: Aspects of prostaglandin function in the lung. *N Engl J Med* 1977; 296:850-855, 910-914
187. Hyman AL, Mathe AA, Lippman HL, et al: Prostaglandins and the lung. *Med Clin North Am* 1981; 65:789-808
188. Carlson LA, Ekelund LG, Oro L: Clinical, metabolic and cardiovascular effects of different prostaglandins in man. *Acta Med Scand* 1970; 188:553-559
189. Kadowitz PJ, Joiner PD, Hyman AL: Effects of prostaglandin E<sub>2</sub> on pulmonary vascular resistance in the intact dog, swine and lamb. *Eur J Pharmacol* 1975; 31:72-75
190. Ishizaki T, Miyabo S, Mifune J, et al: OP-1206, a prostaglandin E<sub>1</sub> derivative: Effects of oral administration to patients with chronic lung disease. *Chest* 1984; 85:382-386
191. Watkins WD, Peterson MB, Crone RK, et al: Prostacyclin and prostaglandin E<sub>1</sub> for severe idiopathic pulmonary artery hypertension (Letter). *Lancet* 1980; 1:1083
192. Mathe AA, Hedqvist P, Holmgren A, et al: Bronchial hyperactivity to prostaglandin F<sub>2</sub> alpha and histamine in patients with asthma. *Br Med J* 1973; 1:193-196
193. Cuthbert MF: Effect on airways resistance of PGE<sub>1</sub> given by aerosol to healthy and asthmatic volunteers. *Br Med J* 1969; 4:723-726
194. Dahlen SE, Hedqvist P, Hammarstrom S, et al: Leukotrienes are potent constrictors of human bronchi. *Nature* 1980; 288:484-486
195. Weiss JW, Drazen JM, Coles N, et al: Bronchoconstrictor effects of leukotriene C in humans. *Science* 1982; 216:196-198
196. Drazen JM, Austen KF, Lewis RA, et al: Comparative airway and vascular activities of leukotrienes C-1 and D in vivo and in vitro. *Proc Natl Acad Sci USA* 1980; 77:4354-4358
197. Bisgaard H: Leukotrienes and prostaglandins in asthma. *Allergy* 1984; 39:413-420
198. Brocklehurst WE: The release of histamine and formation of a slow-reacting substance during anaphylactic shock. *J Physiol (Lond)* 1960; 151:416-435
199. Lewis RA, Austen KF, Drazen JM, et al: Slow-reacting substances of anaphylaxis: Identification of leukotrienes C and D from human and rat sources. *Proc Natl Acad Sci USA* 1980; 77:3710-3714
200. MacGlashan DW, Schleimer RP, Peters SP, et al: Generation of leukotrienes by purified human lung mast cells. *J Clin Invest* 1982; 70:747-751
201. Dahlen SE, Hansson G, Hedqvist P, et al: Allergic challenge of lung tissue from asthmatics elicits bronchial contraction that correlates with the release of leukotriene C<sub>4</sub>, D<sub>4</sub> and E<sub>4</sub>. *Proc Natl Acad Sci USA* 1983; 80:1712-1716
202. Griffin M, Weiss W, Leitch AG, et al: Effects of leukotriene D on the airways in asthma. *N Engl J Med* 1983; 308:436-439
203. Piper PJ, Samhoun MN: The mechanism of action of leukotriene C<sub>4</sub> and D<sub>4</sub> in guinea-pig isolated perfused lung and parenchymal strips of guinea pig lung. *Prostaglandins* 1981; 21:793-803
204. Jones TR, Davis C, Daniel EE: Pharmacological study of the contractile activity of leukotriene C<sub>4</sub> and D<sub>4</sub> on isolated human airway smooth muscle. *Can J Physiol Pharmacol* 1982; 60:638-643
205. Szczeklik A, Gryglewski RJ, Czerniawska-Mysik G: Clinical patterns of hypersensitivity to nonsteroidal anti-inflammatory drugs and their pathogenesis. *J Allergy Clin Immunol* 1977; 60:276-284
206. Hechtman HB, Huval WV, Mathieson MA, et al: Prostaglandin and thromboxane mediation of cardiopulmonary failure. *Surg Clin North Am* 1983; 63:263-283
207. Demling RH: Role of prostaglandins in acute pulmonary microvascular injury. *Ann NY Acad Sci* 1982; 384:517-534
208. Utsunomiya T, Krausz MM, Levine L, et al: Thromboxane mediation of cardiopulmonary effects of embolism. *J Clin Invest* 1982; 70:361-368
209. Klotz TA, Cohn LS, Zipser RD: Urinary excretion of thromboxane B<sub>2</sub> in patients with venous thromboembolic disease. *Chest* 1984; 85:329-335
210. Saldeen T: Clotting, microembolism, and inhibition of fibrinolysis in adult respiratory distress. *Surg Clin North Am* 1983; 63:285-304

211. McKeen CR, Brigham KL, Bowers RE, et al: Pulmonary vascular effects of fat emulsion infusion in unanesthetized sheep—Prevention by indomethacin. *J Clin Invest* 1978; 61:1291-1297
212. Casey LC, Fletcher JR, Zmudka MI, et al: Prevention of endotoxin-induced pulmonary hypertension in primates by the use of a selective thromboxane synthetase inhibitor, OXY 1581. *J Pharmacol Exp Ther* 1982; 222:441-446
213. Jacobs ER, Bone RC: Mediators of septic lung injury. *Med Clin North Am* 1983; 67:701-715
214. Cook JA, Wise WC, Halushka PV: Elevated thromboxane levels in the rat during endotoxic shock: Protective effects of imidazole, 13-azaprostanoic acid, or essential fatty acid deficiency. *J Clin Invest* 1980; 65:227-230
215. Reines HD, Halushka PV, Cook JA, et al: Plasma thromboxane concentrations are raised in patients dying with septic shock. *Lancet* 1982; 2:174-175
216. Bult H, Herman AG: Prostaglandins and circulatory shock. In Herman AG, Vanhoutte PM, Denolin H, et al (Eds): *Cardiovascular Pharmacology of the Prostaglandins*. New York, Raven Press, 1982, pp 327-345
217. Fletcher JR: Prostaglandin synthetase inhibitors in endotoxin or septic shock—A review. *Adv Shock Res* 1983; 10:9-14
218. Clark JG, Kostal KM, Marino BA: Bleomycin-induced pulmonary fibrosis in hamsters: An alveolar macrophage product increases fibroblast prostaglandin  $E_2$  and cyclic adenosine monophosphate and suppresses fibroblast proliferation and collagen production. *J Clin Invest* 1983; 72:2082-2091
219. Goldstein RH, Nukkem J, Glassroth J, et al: Influence of asbestos fibers on collagen and prostaglandin production in fibroblast and macrophage co-cultures. *J Lab Clin Med* 1982; 100:778-785
220. Karim SMM, Filshie GM: Therapeutic abortion using  $PGF_{2\alpha}$ . *Lancet* 1970; 1:157-159
221. Horton EW, Main IHM, Thompson CJ, et al: Effect of orally administered prostaglandin  $E_2$  on gastric secretion and gastrointestinal motility in man. *Gut* 1968; 9:655-658
222. Matuchansky C, Bernier JJ: Effect of  $PGE_1$  on glucose, water and electrolyte absorption in the human jejunum. *Gastroenterology* 1973; 64:1111-1118
223. Cummings JH, Newman A, Misiewicz JJ, et al: Effect of intravenous prostaglandin  $F_{2\alpha}$  on small intestinal function in man. *Nature* 1973; 243:169-171
224. Korman ST, Berant M, Alon U: Review: Prostaglandins in diarrheal states. *Isr J Med Sci* 1981; 17:1109-1113
225. Rachmilewitz DR: Prostaglandins and diarrhea. *Dig Dis Sci* 1980; 25:897-899
226. Rask-Madsen J, Bukhave K, Bytzer P, et al: Prostaglandins in the gastrointestinal tract. *Acta Med Scand* 1984; 685(suppl):30-46
227. Wilson DE, Kaymakcalan H: Prostaglandins: Gastrointestinal effects and peptic ulcer disease. *Med Clin North Am* 1981; 65:773-787
228. Robert A: Prostaglandins in gastrointestinal disease. In Karim SMM (Ed): *Practical Applications of Prostaglandins and Their Synthesis Inhibitors*. Baltimore, University Park Press, 1979, pp 89-102
229. Kinberg DV, Field M, Gershon E, et al: Effects of prostaglandins and cholera enterotoxin on intestinal mucosal cyclic AMP accumulation. *J Clin Invest* 1974; 53:941-949
230. Bukhave K, Rask-Madsen J: Saturation kinetics applied to in vitro effects of low prostaglandin  $E_2$  and  $F_{2\alpha}$  concentrations on ion transport across human jejunal mucosa. *Gastroenterology* 1980; 78:32-42
231. Gots RE, Formai SB, Giannella RA: Indomethacin inhibition of Salmonella typhimurium, Shigella flexneri, and cholera-mediated rabbit ileal secretion. *J Infect Dis* 1974; 130:280-284
232. Metz SA, McRae JR, Robertson RP: Prostaglandins as mediators of paraneoplastic syndromes: Review and up-date. *Metabolism* 1981; 30:299-316
233. Roberts LJ II, Sweetman BJ, Maas RL, et al: Clinical application of PG and TX metabolite quantification. *Prog Lipid Res* 1981; 20:117-121
234. Ligumsky MF, Karmeli F, Sharon P, et al: Enhanced thromboxane  $A_2$  and prostacyclin production by cultured rectal mucosa in ulcerative colitis and its inhibition by steroids and sulfasalazine. *Gastroenterology* 1981; 81:444-449
235. Rampton DS, Hawkey CJ: Prostaglandins and ulcerative colitis (Review). *Gut* 1984; 25:1399-1413
236. Sharon P, Ligumsky M, Rachmilewitz D, et al: Role of prostaglandins in ulcerative colitis—Enhanced production during active disease and inhibition by sulfasalazine. *Gastroenterology* 1978; 75:638-640
237. Craven PA, Saito R, DeFubertis FR: Role of local prostaglandin synthesis in the modulation of proliferative activity of rat colonic epithelium. *J Clin Invest* 1983; 72:1365-1375
238. Campieri M, Lanfranchi GA, Bazzocchi G, et al: Prostaglandins, indomethacin, and ulcerative colitis (Letter). *Gastroenterology* 1980; 78:193
239. Rampton DS, Sladen GE: Prostaglandin synthesis inhibitors in ulcerative colitis: Flurbiprofen compared with conventional treatment. *Prostaglandins* 1981; 21:417-425
240. Campieri M, Lanfranchi GA, Bazzocchi G, et al: Salicylate other than 5-aminosalicylic acid ineffective in ulcerative colitis (Letter). *Lancet* 1978; 2:993
241. Goldin E, Rachmilewitz D: Prostanoids cytoprotection for maintaining remission in ulcerative colitis—Failure of 15(R), 15-methylprostaglandin  $E_2$ . *Dig Dis Sci* 1983; 28:807-811
242. Sharon P, Stenson WF: Enhanced synthesis of leukotriene  $B_4$  by colonic mucosa in inflammatory bowel disease. *Gastroenterology* 1984; 86:453-460
243. Sharon P, Stenson WF: Metabolism of arachidonic acid in acetic acid colitis in rats—Similarity to human inflammatory bowel disease. *Gastroenterology* 1985; 88:55-63
244. Musch MW, Miller RJ, Field M, et al: Stimulation of colonic secretion by lipoxygenase metabolites of arachidonic acid. *Science* 1982; 217:1255-1256
245. Stenson WF, Lobos E: Sulfasalazine inhibits the synthesis of chemotactic lipids by neutrophils. *J Clin Invest* 1982; 69:494-497
246. Robert A, Nezamiste JE, Phillips JP: Inhibition of gastric secretion by prostaglandins. *Am J Dig Dis* 1967; 72:1073-1076
247. Wada T, Ishizawa M: Effects of prostaglandin on the function of the gastric secretion. *Jpn J Clin Med* 1970; 28:2465-2468
248. Wilson DE, Phillips C, Levine RA: Inhibition of gastric secretion in man by prostaglandin  $A_1$ . *Gastroenterology* 1971; 61:201-206
249. Miller TA: Protective effects of prostaglandins against gastric mucosal damage: Current knowledge and proposed mechanisms. *Am J Physiol* 1983; 245:G601-G623
250. Johansson C, Bergström S: Prostaglandins and protection of the gastroduodenal mucosa. *Scand J Gastroenterol* 1982; 17(suppl 77):47-62
251. Check WA: Prostaglandins being tested for gastrointestinal ulcers, bleeding. *Clin Pharm* 1984; 3:563-568
252. Goyal RK: Deleterious effects of prostaglandins on esophageal mucosa (Editorial). *Gastroenterology* 1980; 78:1085-1087
253. Northway MG, Castell DO: Do prostaglandins cause gastrointestinal mucosal injury? *Dig Dis Sci* 1981; 26:453-456
254. Hollander D, Tarnawski A, Gergely M, et al: Sucralfate protection of the gastric mucosa against ethanol induced injury: A prostaglandin-mediated process? *Scand J Gastroenterol* 1984; 19(suppl):97-102
255. Thornell E: Mechanisms in the development of acute cholecystitis and biliary pain: A study of the role of prostaglandins and effects of indomethacin. *Scand J Gastroenterol* 1982; 17(suppl 76):1-31
256. Fiedler L:  $PGF_{2\alpha}$ —A new therapy for paralytic ileus? *Adv Prostaglandin Thromboxane Res* 1980; 8:1609-1610
257. Robertson RP: Prostaglandins, glucose homeostasis, and diabetes mellitus. *Med Clin North Am* 1981; 65:759-771
258. Robertson RP: Prostaglandins, glucose homeostasis, and diabetes mellitus. *Annu Rev Med* 1983; 34:1-12
259. Robertson RP: Arachidonic acid and metabolic disorders. *Spec Top Endocrinol Metab* 1983; 5:55-81
260. Metz S, VanRollins M, Strife R, et al: Lipoxygenase pathway in islet endocrine cells—Oxidative metabolism of arachidonic acid promotes insulin release. *J Clin Invest* 1983; 71:1191-1205
261. Voelkel EF, Tashjian AH Jr, Franklin R, et al: Hypercalcemia and tumor-prostaglandins: The  $VX_2$  carcinoma model in the rabbit. *Metabolism* 1975; 24:973-986
262. Doppelt SH, Slovick DM, Neer RM, et al: Gut-mediated hypercalcemia in rabbits bearing  $VX_2$  carcinoma: New mechanism for tumor-induced hypercalcemia. *Proc Natl Acad Sci USA* 1982; 79:640-644
263. Seyberth HW, Segre GV, Morgan JL, et al: Prostaglandins as mediators of hypercalcemia associated with certain types of cancer. *N Engl J Med* 1975; 293:1278-1283
264. Robertson RP, Baylink DJ, Metz SA, et al: Plasma prostaglandin  $E$  in patients with cancer with and without hypercalcemia. *J Clin Endocrinol Metab* 1976; 43:1330-1335
265. Demers LM, Allegra JC, Harvey HA, et al: Plasma prostaglandins in hypercalcemic patients with neoplastic disease. *Cancer* 1977; 39:1559-1562
266. Minkin C, Fredericks RS, Pokress S, et al: Bone resorption and humoral hypercalcemia of malignancy: Stimulation of bone resorption in vitro by tumor extracts is inhibited by prostaglandin synthesis inhibitors. *J Clin Endocrinol Metab* 1981; 53:941-947
267. Mundy GR, Ibbotson KJ, D'Souza SM, et al: The hypercalcemia of cancer: Clinical implications and pathogenic mechanism. *N Engl J Med* 1984; 310:1718-1727
268. Goodwin JS: Prostaglandins and host defense in cancer. *Med Clin North Am* 1981; 65:829-844
269. Levine L: Arachidonic acid transformation and tumor production. *Adv Cancer Res* 1981; 35:49-79
270. Honn KV, Menter D, Cavanaugh PG, et al: A review of prostaglandins and the treatment of tumor metastasis. *Acta Clin Belg* 1983; 38:53-67
271. Tisdale MJ: Role of prostaglandins in metastatic dissemination of cancer: Minireview on cancer research. *Exp Cell Biol* 1983; 51:250-256
272. Bockman RS: Prostaglandins in cancer: A review. *Cancer Invest* 1983; 1:485-493
273. Bennett A: Prostaglandins and cancer. In Karim SMM (Ed): *Practical Applications of Prostaglandins and Their Synthesis Inhibitors*. Baltimore, University Park Press, 1979, pp 149-188